

Lamivudine (3TC, Epivir)

Updated: September 30, 2025

Reviewed: September 30, 2025

Formulations	
<p>Pediatric Oral Solution</p> <ul style="list-style-type: none"> • [Epivir] 10 mg/mL • [Epivir HBV]^a 5 mg/mL <p>Tablets</p> <ul style="list-style-type: none"> • [Epivir] 150 mg (scored) and 300 mg • [Epivir HBV]^a 100 mg <p>Generic Formulations</p> <ul style="list-style-type: none"> • 100-mg, 150-mg, and 300-mg tablets <p>Fixed-Dose Combination (FDC) Tablets</p> <ul style="list-style-type: none"> • [Cimduo] Lamivudine 300 mg/tenofovir disoproxil fumarate 300 mg • [Lamivudine 150 mg/zidovudine 300 mg • [Delstrigo] Doravirine 100 mg/lamivudine 300 mg/tenofovir disoproxil fumarate 300 mg • [Dovato] Dolutegravir 50 mg/lamivudine 300 mg • Abacavir 600 mg/lamivudine 300 mg • [Symfi] Efavirenz 600 mg/lamivudine 300 mg/tenofovir disoproxil fumarate 300 mg • Efavirenz 400 mg/lamivudine 300 mg/tenofovir disoproxil fumarate 300 mg • [Temixys] Lamivudine 300 mg/tenofovir disoproxil fumarate 300 mg • [Triumeq] Abacavir 600 mg/dolutegravir 50 mg/lamivudine 300 mg • [Triumeq PD] Abacavir 60 mg/dolutegravir 5 mg/lamivudine 30 mg <p>When using FDC tablets, refer to other sections of Appendix A. Pediatric Antiretroviral Drug Information for information about the individual components of the FDC. See also Appendix A, Table 2. Antiretroviral Fixed-Dose Combination Tablets: Minimum Body Weights and Considerations for Use in Children and Adolescents.</p> <p>For additional information, see Drugs@FDA or DailyMed.</p>	
Dosing Recommendations	Selected Adverse Events
<p>Note: See Antiretroviral Management of Infants With <i>In Utero</i>, Intrapartum, or Breastfeeding Exposure to HIV and Table 13.1. Drug Dosing Recommendations for Antiretroviral Prophylaxis and Presumptive HIV Therapy in Infants With <i>In Utero</i> or Intrapartum Exposure to HIV for information about using lamivudine (3TC) to prevent perinatal HIV transmission.</p>	<ul style="list-style-type: none"> • Headache
	<p>Special Instructions</p> <ul style="list-style-type: none"> • 3TC and coformulated tablets can be given with and without food. • Store 3TC oral solution at room temperature.

Neonate (≥32 Weeks Gestation at Birth) and Infant (Birth to <4 Weeks) Dose**Oral Solution**

- 3TC 2 mg/kg twice daily

Infant and Child Dose

- Once-daily dosing of the 3TC oral solution is **not recommended** when initiating 3TC oral solution in infants and young children. Patients can be transitioned to once-daily treatment with the oral solution when they have been stable on twice-daily treatment for 36 weeks and are aged ≥3 years. Please see the note below and refer to the text for more detail.

Aged ≥4 Weeks to <3 Months

- 3TC 4 mg/kg twice daily of the oral solution

Aged ≥3 Months to <3 Years

- 3TC 5 mg/kg twice daily of the oral solution (maximum 150 mg per dose)

Aged ≥3 Years

- 3TC 5 mg/kg twice daily of the oral solution (maximum 150 mg per dose); or
- 3TC 10 mg/kg once daily of the oral solution (maximum 300 mg per dose)

Weight-Band Dosing for the 10-mg/mL 3TC Oral Solution in Children Weighing ≥3 kg and Aged ≥4 Weeks

Weight	Twice-Daily Dose, AM	Twice-Daily Dose, PM
3 kg to <6 kg	3 mL	3 mL
6 kg to <10 kg	4 mL	4 mL
10 kg to <14 kg	6 mL	6 mL

Note: Weight-band dosing in the table above is derived from World Health Organization recommendations. These doses result in significantly higher 3TC exposure than mg per kg doses, especially for young infants. However, given that 3TC has a generally favorable safety profile, these doses are thought to be safe and can be used as soon as infants weigh ≥3 kg and are aged ≥4 weeks.

Sources: Tremoulet AH, Nikanjam M, Cressey TR, et al. Developmental pharmacokinetic changes of lamivudine in infants and children. *J Clin Pharmacol.* 2012;52(12):1824-1832. Available at: <https://pubmed.ncbi.nlm.nih.gov/22180560>. World Health Organization. Annex 3. Dosages of ARV drugs for adults and adolescents. 2018. Available at: https://cdn.who.int/media/docs/default-source/hq-hiv-hepatitis-and-stis-library/arv_guidelines-2018-annex3a.pdf?sfvrsn=1318b82d_3.

- For abacavir (ABC)/dolutegravir (DTG)/3TC dispersible tablets, fully disperse them in 20 mL of drinking water in the supplied cup and swirl the suspension so that no lumps remain. After full dispersion and within 30 minutes of mixing, administer the oral suspension. Rinse the dosing cup with a small amount of water, and give this additional water to the child to ensure that the child takes the full dose and no medication remains in the dosing cup. ABC/DTG/3TC dispersible tablets should not be swallowed whole, chewed, cut, or crushed.
- Screen patients for hepatitis B virus (HBV) infection before using 3TC or FDC tablets that contain 3TC. Severe acute exacerbations of HBV can occur after discontinuation of 3TC. Hepatic function and HBV viral load should be monitored for several months after patients with HBV infection stop taking 3TC. Patients with HBV/HIV coinfection who receive Dovato will require additional treatment for chronic HBV infection.
- For any FDC tablet containing ABC, test patients for the HLA-B*5701 allele before starting therapy to predict the risk of hypersensitivity reactions. Patients who test positive for the HLA-B*5701 allele should not be given an ABC-containing FDC. Patients with no prior HLA-B*5701 testing who are tolerating an ABC-containing regimen do not need to be tested. See [Abacavir](#).

Metabolism/Elimination**3TC Dosing in Patients With Hepatic Impairment**

- No change in 3TC dosing is required for patients with hepatic impairment.
- FDC tablets containing ABC or ZDV should not be used in patients who have impaired hepatic function.
- Symfi (EFV 600 mg/3TC/TDF) and the FDC containing EFV 400 mg/3TC/TDF should be used with caution in patients with hepatic impairment; these FDCs are not recommended for use in moderate or severe hepatic impairment.

Weighting ≥14 kg and Able to Swallow Tablets

- Weight-band dosing (see table below; dose is approximately 3TC 5 mg/kg per day twice daily or 3TC 10 mg/kg once daily)
- The scored tablet is the preferred formulation for pediatric patients weighing ≥14 kg who can swallow a tablet.

Weight-Band Dosing for the Scored 150-mg 3TC Tablet in Children Weighing ≥14 kg

Weight	Twice-Daily Dose, AM	Twice-Daily Dose, PM	Once-Daily Dose
14 kg to <20 kg	½ tablet (75 mg)	½ tablet (75 mg)	1 tablet (150 mg)
≥20 kg to <25 kg	½ tablet (75 mg)	1 tablet (150 mg)	1½ tablets (225 mg)
≥25 kg	1 tablet (150 mg)	1 tablet (150 mg)	2 tablets (300 mg)

Note: The Panel on Antiretroviral Therapy and Medical Management of Children Living With HIV (the Panel) supports switching from twice-daily dosing to once-daily dosing of 3TC (using the oral solution or tablets) in children aged ≥3 years who have been clinically stable for 36 weeks with undetectable viral loads and stable CD4 T lymphocyte cell counts. Clinicians should choose a once-daily regimen using the once-daily dose of 3TC indicated above (approximately 3TC 10 mg/kg, with a maximum of 3TC 300 mg once daily).

Child and Adolescent (Weighting ≥25 kg) and Adult Dose

- 3TC 150 mg twice daily; or
- 3TC 300 mg once daily

[Cimduo] 3TC/Tenofovir Disoproxil Fumarate (TDF)

Child and Adolescent (Weighting >35 kg) and Adult Dose

- One tablet once daily

3TC/Zidovudine (ZDV)

Child and Adolescent (Weighting ≥30 kg) and Adult Dose

- One tablet twice daily

[Delstrigo] Doravirine/3TC/TDF

Child and Adolescent (Weighting ≥35 kg) and Adult Dose

- One tablet once daily in antiretroviral (ARV)-naïve patients and ARV-experienced patients who have been virologically suppressed (HIV RNA <50 copies/mL) on a stable ARV regimen with no history of treatment failure and no known mutations associated with resistance to the individual components of Delstrigo

- Delstrigo and Dovato do not require dose adjustment in mild or moderate hepatic impairment but have not been studied and, therefore, are not recommended in patients with severe hepatic impairment.

3TC Dosing in Patients With Renal Impairment

- Dose adjustment of 3TC is required for patients with renal insufficiency.
- Do not use FDC tablets containing 3TC in patients with creatinine clearance <30 mL/min or patients on dialysis because the doses of 3TC cannot be adjusted. Data from the FDC DTG/3TC (Dovato) suggest that patients with a sustained creatinine clearance of 30–49 mL/min may experience a higher 3TC exposure and should be monitored for hematologic toxicities and potential FDC discontinuation and subsequent adjustment of the treatment regimen. See package inserts for additional information.

[Dovato] Dolutegravir (DTG)/3TC

Adolescent (Weighing ≥ 25 kg and aged ≥ 12 years) and Adult Dose

- One tablet once daily with or without food as a complete ARV regimen in antiretroviral therapy (ART)—naïve adults and adolescents with no known mutations associated with resistance to the individual components of Dovato.
- The combination tablet DTG/3TC (Dovato) is approved by the U.S. Food and Drug Administration (FDA) for adolescents weighing ≥ 25 kg and aged ≥ 12 years but is not approved for use in children aged < 12 years. However, it could be used as part of a three-drug regimen in patients who meet the minimum body weight requirements for each component of the drug.

Abacavir (ABC)/3TC

Child and Adolescent (Weighing ≥ 25 kg) and Adult Dose

- One tablet once daily

[Symfi] Efavirenz (EFV) 600 mg/3TC/TDF

Child and Adolescent (Weighing ≥ 40 kg) and Adult Dose

- One tablet once daily on an empty stomach

EFV 400 mg/3TC/TDF

Child and Adolescent (Weighing ≥ 35 kg) and Adult Dose

- One tablet once daily on an empty stomach
- The FDC containing 400mg EFV/3TC/TDF has not been studied in children (sexual maturity ratings [SMRs] 1–3), and major interindividual variability in EFV plasma concentrations has been found in pediatric patients in a multiethnic setting. The 400-mg dose of EFV may be too low in children or adolescents with SMRs 1 to 3 who weigh ≥ 40 kg. The use of therapeutic drug monitoring is suggested by some Panel members when this FDC is used in pediatric patients who weigh ≥ 40 kg (see the [Efavirenz](#) section for more information).

[Temixys] 3TC/TDF

Child and Adolescent (Weighing ≥ 35 kg) and Adult Dose

- One tablet once daily

[Triumeq PD] ABC/DTG/3TC

Child Weighing ≥ 6 kg to < 25 kg and Aged ≥ 3 Months

- Dispersible Triumeq PD tablets are FDA approved for children weighing ≥ 6 to < 25 kg. Triumeq PD is not recommended for children weighing ≥ 25 kg who are eligible for adult Triumeq dosing.
- Administer the appropriate number of tablets for a child's weight once daily, dispersed in 20 mL of water. See Special Instructions. Triumeq PD tablets should not be swallowed whole, chewed, cut, or crushed.

Weight-Band Dosing of Triumeq PD Tablets for Children Weighing ≥6 kg

Weight	Recommended Daily Dose	Number of Triumeq PD Tablets
6 kg to <10 kg	ABC 180 mg DTG 15 mg 3TC 90 mg	3
10 kg to <14 kg	ABC 240 mg DTG 20 mg 3TC 120 mg	4
14 kg to <20 kg	ABC 300 mg DTG 25 mg 3TC 150 mg	5
20 kg to <25 kg	ABC 360 mg DTG 30 mg 3TC 180 mg	6
≥25 kg	Use Triumeq (see below).	

- For use in children who are ARV-naïve or ARV-experienced (but integrase strand transfer inhibitor [INSTI]-naïve) and who are not being treated with uridine diphosphate glucuronosyltransferase 1A1 (UGT1A1) or cytochrome P450 (CYP) 3A inducers

[Triumeq] ABC/DTG/3TC

Child and Adolescent (Weighing ≥25 kg) and Adult Dose

- One tablet once daily
- This FDC tablet can be used in patients who are ART-naïve or ART-experienced (but INSTI-naïve) and who are not being treated with UGT1A1 or CYP3A inducers.

^a EpiVir HBV oral solution and tablets contain a lower amount of 3TC than EpiVir oral solution and tablets. The amount of 3TC in the EpiVir HBV solution and tablet was based on dosing for treatment of HBV infection in people without HIV coinfection. Patients with HIV who are taking EpiVir HBV as part of their ARV regimen should receive the appropriate amount of oral solution or the appropriate number of tablets to achieve the higher doses of 3TC that are used to treat HIV.

Drug Interactions

Additional information about drug interactions is available in the [Adult and Adolescent Antiretroviral Guidelines](#) and the [HIV Drug Interaction Checker](#).

- Drugs that decrease renal function could decrease clearance of lamivudine (3TC).
- Do not use** 3TC in combination with emtricitabine (FTC) because these drugs have similar resistance profiles and using them together offers no additional benefit.¹ **Do not use** 3TC with fixed-dose combination (FDC) medications that contain 3TC or FTC. Please see [Appendix A, Table 1. Antiretrovirals Available in Fixed-Dose Combination Tablets or as a Co-packaged Formulation, by Drug Class](#) and refer to other sections of the [Drug Appendix](#) for drug interaction information about each individual component of FDC tablets.

Major Toxicities

- *More common:* Headache, nausea
- *Less common (more severe):* Peripheral neuropathy, lipodystrophy/lipoatrophy
- *Rare:* Increased levels of liver enzymes. Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported.

Resistance

The International Antiviral Society–USA maintains a list of [HIV drug resistance mutations](#), and the [Stanford University HIV Drug Resistance Database](#) offers a discussion of each mutation.

Pediatric Use

Approval

Although 3TC is approved by the U.S. Food and Drug Administration (FDA) for the treatment of children aged ≥ 3 months, both the Panel on Antiretroviral Therapy and Medical Management of Children Living With HIV (the Panel) and the Panel on Treatment of HIV During Pregnancy and Prevention of Perinatal Transmission recommend the use of 3TC from birth based on extensive experience, as well as safety and pharmacokinetic (PK) data in this age group.

Considerations for Use

The efficacy and toxicity of 3TC are equivalent to the efficacy and toxicity of FTC. The oral formulation of FTC has an advantage over the liquid formulation of 3TC because it can be given once daily at antiretroviral (ARV) initiation, whereas the liquid formulation of 3TC needs to be given twice daily at ARV initiation. When pill formulations of 3TC or FTC are used, they can be administered once daily.

Both FTC and 3TC have antiviral activity and efficacy against hepatitis B virus. For a comprehensive review of this topic, see the [Hepatitis B Virus](#) section in the [Pediatric Opportunistic Infection Guidelines](#).

Comparative Clinical Trials

Investigators studying the AIDS Therapy Evaluation in the Netherlands (ATHENA) cohort compared the efficacy of tenofovir disoproxil fumarate (TDF) plus FTC to TDF plus 3TC when these drugs were administered with a ritonavir-boosted protease inhibitor (darunavir, atazanavir, or lopinavir) in ART-naïve patients.² The adjusted hazard ratio for the virologic failure of 3TC-containing regimens compared to FTC-containing regimens within 240 weeks of starting therapy was 1.15 (95% confidence interval, 0.58–2.27). These regimens had no difference in time to virologic suppression during the first 48 weeks of therapy or time to virologic failure after attaining suppression. In a Swiss cohort, Yang et al. found a potential difference in efficacy between FTC and 3TC; however, the difference disappeared after adjusting for pill burden. Current evidence suggests that FTC and 3TC have equivalent efficacy and toxicity in ARV-naïve patients.³

Efficacy

3TC has been studied in children with HIV both alone and in combination with other ARV drugs. Extensive data have demonstrated the safety of 3TC and have shown that this drug is associated

with clinical improvement and virologic response. It is commonly used in children with HIV as a component of a dual nucleoside reverse transcriptase inhibitor (NRTI) backbone.⁴⁻¹² In one study that evaluated the efficacy of NRTI background components, the combination of 3TC plus abacavir (ABC) was superior to zidovudine (ZDV) plus 3TC or ZDV plus ABC in achieving long-term virologic efficacy.¹³

Pharmacokinetics in Infants

Because of its safety profile and availability in a liquid formulation, 3TC has been given to infants during the first 6 weeks of life starting at a dose of 2 mg/kg every 12 hours before age 4 weeks.⁹ A population PK analysis of infants who received 3TC affirms that adjusting the dose from 3TC 2 mg/kg to 3TC 4 mg/kg every 12 hours at age 4 weeks provides optimal 3TC exposure for infants with normal maturation of renal function.¹⁴ For infants, the World Health Organization (WHO) weight-band dosing (which is up to five times higher than the FDA-approved dose) results in greater plasma concentrations than the 3TC 2 mg/kg dose.¹⁵ In HIV Prevention Trials Network 040, 3TC was administered as a component of a three-drug regimen to prevent perinatal transmission during the first 2 weeks of life. For 2 weeks, all infants weighing >2,000 g received 3TC 6 mg twice daily, and infants weighing ≤2,000 g received 3TC 4 mg twice daily. These doses resulted in 3TC exposure that was similar to the exposure seen in infants who received the standard twice-daily dosing schedule of 3TC 2 mg/kg per dose for neonates.¹⁶

In preterm infants, PK modeling supports the use of 3TC for infants. For those with gestational age at birth of ≥24 to <30 weeks, a dose of 2 mg/kg twice daily from birth until weighing 3 kg is supported. For infants with gestational age at birth of ≥30 to <37 weeks, 2 mg/kg twice daily for the first 4 weeks of life followed by 4 mg/kg twice daily is supported. This proposed 3TC dosing has been endorsed by the WHO Pediatric Antiretroviral Working Group but has not yet been clinically validated in preterm infants.¹⁷

Pharmacokinetics of Liquid Versus Tablet Preparations

The PK of 3TC have been studied after either single or repeat doses in 210 pediatric subjects. Pediatric subjects who received 3TC oral solution according to the recommended dose regimen achieved plasma concentrations of 3TC that were approximately 25percent lower compared with those of adults with HIV who received the oral solution. Pediatric subjects who received 3TC tablets achieved plasma concentrations that were comparable to or slightly higher than those observed in adults who received tablets. In pediatric subjects, the relative bioavailability of 3TC oral solution is approximately 40% lower than the relative bioavailability of tablets that contain 3TC, despite no difference in the bioavailability of these two formulations among adults. The mechanisms for the diminished relative bioavailability of 3TC oral solution are unknown,¹⁸ but results from a study in adults that compared the PK of 3TC oral solution administered either alone or with increasing concentrations of sorbitol indicate that sorbitol decreases the total exposure of 3TC oral solution.¹⁹ Sorbitol is a component of several ARV solutions, including ABC, as well as common over-the-counter medications that may be used in infants and young children; this may explain the PK discrepancy between the oral solution and tablet formulations. Modeling of PK data in pediatric patients suggests that increasing the oral solution dose to 3TC 5 mg/kg per dose twice daily or 3TC 10 mg/kg per dose once daily (with a maximum of 3TC 300 mg administered daily) in children aged ≥3 months would provide exposures similar to those seen in adult patients who received tablet formulations. However, these models used PK data derived from studies that did not use 3TC liquid formulation, so modeling may not predict exposures for 3TC oral solution, especially when used with liquid ABC. The Panel **does not recommend** using a once-daily dose of 3TC until a child is aged ≥3 years. After 3 years of age, switching to once-daily dosing with the liquid formulation could be considered when harmonizing with other ARV drugs administered once daily, such as ABC and dolutegravir (DTG).

Dosing Considerations—Once-Daily Versus Twice-Daily Administration

The standard adult dose for 3TC is 300 mg once daily, but data are lacking on once-daily administration of 3TC in children. Population PK data indicate that once-daily dosing of 3TC 8 mg/kg leads to area under the curve over 24 hours (AUC_{0-24h}) values that are similar to those seen in patients taking 3TC 4 mg/kg twice daily, but minimum blood plasma concentration (C_{min}) values are significantly lower and maximum blood plasma concentration (C_{max}) values are significantly higher in children aged 1 year to 18 years.²⁰ Intensive PK of once-daily versus twice-daily dosing of 3TC were evaluated in children with HIV aged 2 to 13 years in the PENTA (Paediatric European Network for Treatment of AIDS) 13 trial⁴ and in children aged 3 months to 36 months in the PENTA 15 trial.²¹ Both the PENTA 13 and PENTA 15 trials used a crossover design with doses of 3TC 8 mg/kg once daily or 3TC 4 mg/kg twice daily. AUC_{0-24h} and clearance values were similar between these two dosing schedules, and most children maintained an undetectable HIV RNA value after the switch. An ARROW (AntiRetroviral Research fOr Watoto) trial PK study of 41 children aged 3 to 12 years (median age 7.6 years) in Uganda who were stable on twice-daily 3TC also showed equivalent AUC_{0-24h} and good clinical outcomes (defined by a low disease stage and a high CD4 T lymphocyte [CD4] cell count) after switching to once-daily 3TC. Median follow-up time during this study was 1.15 years.²² The larger ARROW trial was a randomized, noninferiority trial that investigated once-daily versus twice-daily doses of 3TC in more than 600 pediatric patients who had initiated therapy with twice-daily 3TC and who had been receiving therapy for ≥ 36 weeks. Median follow-up time during the study was 114 weeks. Rates of plasma HIV RNA suppression and adverse event profiles for once-daily 3TC were similar to (and statistically non-inferior to) those of twice-daily 3TC.²³

All four of the studies discussed above enrolled patients who had low plasma HIV RNA or who were clinically stable on twice-daily 3TC before switching to once-daily dosing. Therefore, the Panel supports switching from twice-daily to once-daily dosing of 3TC in children aged ≥ 3 years who have been clinically stable for 36 weeks with an undetectable viral load and stable CD4 count. Clinicians should use a 10 mg/kg per dose of 3TC oral solution or a weight-based dose of 3TC tablets (neither exceeding 3TC 300 mg) as part of a once-daily regimen.²⁴ More long-term clinical trials with viral efficacy endpoints are needed to confirm that once-daily dosing of 3TC can be used effectively as part of an initial ARV regimen in children.

3TC undergoes intracellular metabolism to reach its active form, 3TC triphosphate. In adolescents, the mean half-life of intracellular 3TC triphosphate (17.7 hours) is considerably longer than that of unphosphorylated 3TC in plasma (1.5–2 hours). Intracellular concentrations of 3TC triphosphate are equivalent whether 3TC is given once daily or twice daily in adults and adolescents. This supports a recommendation for once-daily 3TC dosing based on FDA recommendations.^{25,26}

Weight-Band Dosing

Weight-band dosing recommendations for 3TC have been developed for children weighing ≥ 3 kg and receiving either the 10-mg/mL oral solution or the 150-mg scored tablets.²⁷⁻²⁹

Recent data from the IMPAACT 2019 clinical trial of dispersible and immediate-release ABC/DTG/3TC tablets (ABC 60 mg, DTG 5 mg, and 3TC 30 mg dispersed in 15–20 mL of water) in children with HIV has confirmed the FDA-approved dosing in infants and children weighing 6 to <25 kg. ABC/DTG/3TC dispersible FDC dosing was confirmed based on PK and safety data in each weight-band at the originally selected dosing, which aligned with WHO weight-band dosing for the individual ARV agents. Follow-up through 24 weeks confirmed the safety, tolerability, and virological efficacy of both formulations.³⁰ Post hoc population pharmacokinetic modeling of IMPAACT 2019 data further supports weight-band dosing of ABC/3TC/DTG FDC in children weighing ≥ 6 kg.³¹

References

1. Anderson PL, Lamba J, Aquilante CL, et al. Pharmacogenetic characteristics of indinavir, zidovudine, and lamivudine therapy in HIV-infected adults: a pilot study. *J Acquir Immune Defic Syndr*. 2006;42(4):441-449. Available at: <https://pubmed.ncbi.nlm.nih.gov/16791115>.
2. Rokx C, Gras L, van de Vijver D, et al. Virological responses to lamivudine or emtricitabine when combined with tenofovir and a protease inhibitor in treatment-naïve HIV-1-infected patients in the Dutch AIDS Therapy Evaluation in the Netherlands (ATHENA) cohort. *HIV Med*. 2016;17(8):571-580. Available at: <https://pubmed.ncbi.nlm.nih.gov/26842457>.
3. Yang WL, Kouyos RD, Scherrer AU, et al. Assessing efficacy of different nucleos(t)ide backbones in NNRTI-containing regimens in the Swiss HIV Cohort Study. *J Antimicrob Chemother*. 2015;70(12):3323-3331. Available at: <https://pubmed.ncbi.nlm.nih.gov/26362944>.
4. Bergshoeff A, Burger D, Verweij C, et al. Plasma pharmacokinetics of once- versus twice-daily lamivudine and abacavir: simplification of combination treatment in HIV-1-infected children (PENTA-13). *Antivir Ther*. 2005;10(2):239-246. Available at: <https://pubmed.ncbi.nlm.nih.gov/15865218>.
5. Chadwick EG, Rodman JH, Britto P, et al. Ritonavir-based highly active antiretroviral therapy in human immunodeficiency virus type 1-infected infants younger than 24 months of age. *Pediatr Infect Dis J*. 2005;24(9):793-800. Available at: <https://pubmed.ncbi.nlm.nih.gov/16148846>.
6. Chaix ML, Rouet F, Kouakoussui KA, et al. Genotypic human immunodeficiency virus type 1 drug resistance in highly active antiretroviral therapy-treated children in Abidjan, Cote d'Ivoire. *Pediatr Infect Dis J*. 2005;24(12):1072-1076. Available at: <https://pubmed.ncbi.nlm.nih.gov/16371868>.
7. Krogstad P, Lee S, Johnson G, et al. Nucleoside-analogue reverse-transcriptase inhibitors plus nevirapine, nelfinavir, or ritonavir for pretreated children infected with human immunodeficiency virus type 1. *Clin Infect Dis*. 2002;34(7):991-1001. Available at: <https://pubmed.ncbi.nlm.nih.gov/11880966>.
8. LePrevost M, Green H, Flynn J, et al. Adherence and acceptability of once daily lamivudine and abacavir in human immunodeficiency virus type-1 infected children. *Pediatr Infect Dis J*. 2006;25(6):533-537. Available at: <https://pubmed.ncbi.nlm.nih.gov/16732152>.
9. Mirochnick M, Stek A, Acevedo M, et al. Safety and pharmacokinetics of nelfinavir coadministered with zidovudine and lamivudine in infants during the first 6 weeks of life. *J Acquir Immune Defic Syndr*. 2005;39(2):189-194. Available at: <https://pubmed.ncbi.nlm.nih.gov/15905735>.
10. Mueller BU, Lewis LL, Yuen GJ, et al. Serum and cerebrospinal fluid pharmacokinetics of intravenous and oral lamivudine in human immunodeficiency virus-infected children. *Antimicrob Agents Chemother*. 1998;42(12):3187-3192. Available at: <https://pubmed.ncbi.nlm.nih.gov/9835513>.
11. Nachman SA, Stanley K, Yogev R, et al. Nucleoside analogs plus ritonavir in stable antiretroviral therapy-experienced HIV-infected children: a randomized controlled trial. Pediatric AIDS clinical trials group 338 study team. *JAMA*. 2000;283(4):492-498. Available at: <https://pubmed.ncbi.nlm.nih.gov/10659875>.

12. Scherpbier HJ, Bekker V, van Leth F, et al. Long-term experience with combination antiretroviral therapy that contains nelfinavir for up to 7 years in a pediatric cohort. *Pediatrics*. 2006;117(3):e528-536. Available at: <https://pubmed.ncbi.nlm.nih.gov/16481448>.
13. Green H, Gibb DM, Walker AS, et al. Lamivudine/abacavir maintains virological superiority over zidovudine/lamivudine and zidovudine/abacavir beyond 5 years in children. *AIDS*. 2007;21(8):947-955. Available at: <https://pubmed.ncbi.nlm.nih.gov/17457088>.
14. Tremoulet AH, Capparelli EV, Patel P, et al. Population pharmacokinetics of lamivudine in human immunodeficiency virus-exposed and -infected infants. *Antimicrob Agents Chemother*. 2007;51(12):4297-4302. Available at: <https://pubmed.ncbi.nlm.nih.gov/17893155>.
15. Mirochnick M, Nielsen-Saines K, Pilotto JH, et al. Nelfinavir and Lamivudine pharmacokinetics during the first two weeks of life. *Pediatr Infect Dis J*. 2011;30(9):769-772. Available at: <https://pubmed.ncbi.nlm.nih.gov/21666540>.
16. Bekker A, Yang J, Wang J, et al. Safety and pharmacokinetics of lopinavir/ritonavir oral solution in preterm and term infants starting before 3 months of age. *Pediatr Infect Dis J*. 2024;43(4):355-360. Available at: <https://pubmed.ncbi.nlm.nih.gov/38190642>.
17. Choi SY, Li F, Florian J, Seo SK. Lamivudine and abacavir clinical summary review. 2014. Available at: <https://www.fda.gov/downloads/Drugs/DevelopmentApprovalProcess/DevelopmentResources/UCM446104.pdf>.
18. Adkison K, Wolstenholme A, Lou Y, et al. Effect of sorbitol on the pharmacokinetic profile of lamivudine oral solution in adults: an open-label, randomized study. *Clin Pharmacol Ther*. 2018;103(3):402-408. Available at: <https://pubmed.ncbi.nlm.nih.gov/29150845>.
19. Bouazza N, Hirt D, Blanche S, et al. Developmental pharmacokinetics of lamivudine in 580 pediatric patients ranging from neonates to adolescents. *Antimicrob Agents Chemother*. 2011;55(7):3498-3504. Available at: <https://pubmed.ncbi.nlm.nih.gov/21576443>.
20. Paediatric European Network for Treatment of AIDS. Pharmacokinetic study of once-daily versus twice-daily abacavir and lamivudine in HIV type-1-infected children aged 3–<36 months. *Antivir Ther*. 2010;15(3):297-305. Available at: <https://pubmed.ncbi.nlm.nih.gov/20516550>.
21. Musiime V, Kendall L, Bakeera-Kitaka S, et al. Pharmacokinetics and acceptability of once-versus twice-daily lamivudine and abacavir in HIV type-1-infected Ugandan children in the ARROW Trial. *Antivir Ther*. 2010;15(8):1115-1124. Available at: <https://pubmed.ncbi.nlm.nih.gov/21149918>.
22. Musiime V, Kasirye P, Naidoo-James B, et al. Once vs. twice-daily abacavir and lamivudine in African children. *AIDS*. 2016;30(11):1761-1770. Available at: <https://pubmed.ncbi.nlm.nih.gov/27064996>.
23. Janssen EJH, Bastiaans DET, Valitalo PAJ, et al. Dose evaluation of lamivudine in human immunodeficiency virus-infected children aged 5 months to 18 years based on a population pharmacokinetic analysis. *Br J Clin Pharmacol*. 2017;83(6):1287-1297. Available at: <https://pubmed.ncbi.nlm.nih.gov/28079918>.

24. Yuen GJ, Lou Y, Bumgarner NF, et al. Equivalent steady-state pharmacokinetics of lamivudine in plasma and lamivudine triphosphate within cells following administration of lamivudine at 300 milligrams once daily and 150 milligrams twice daily. *Antimicrob Agents Chemother*. 2004;48(1):176-182. Available at: <https://pubmed.ncbi.nlm.nih.gov/14693537>.
25. Flynn PM, Rodman J, Lindsey JC, et al. Intracellular pharmacokinetics of once versus twice daily zidovudine and lamivudine in adolescents. *Antimicrob Agents Chemother*. 2007;51(10):3516-3522. Available at: <https://pubmed.ncbi.nlm.nih.gov/17664328>.
26. World Health Organization. Preferred antiretroviral medicines for treating and preventing HIV infection in younger children: report of the WHO Paediatric Antiretroviral Working Group. 2008. Available at: https://apps.who.int/iris/bitstream/handle/10665/43882/9789241596619_eng.pdf;jsessionid=31285037F69DEBBDEE7C08D16ADD48F7?sequence=1.
27. L'Homme RF, Kabamba D, Ewings FM, et al. Nevirapine, stavudine and lamivudine pharmacokinetics in African children on paediatric fixed-dose combination tablets. *AIDS*. 2008;22(5):557-565. Available at: <https://pubmed.ncbi.nlm.nih.gov/18316996>.
28. Brooks KM, Kiser JJ, Ziemba L, et al. Pharmacokinetics, safety, and tolerability of dispersible and immediate-release abacavir, dolutegravir, and lamivudine tablets in children with HIV (IMPAACT 2019): week 24 results of an open-label, multicentre, phase 1-2 dose-confirmation study. *Lancet HIV*. 2023;10(8):e506-e517. Available at: <https://pubmed.ncbi.nlm.nih.gov/37541705>.
29. Chandasana H, van Dijkman SC, Mehta R, et al. Population pharmacokinetic modeling of abacavir/dolutegravir/lamivudine to support a fixed-dose combination in children with HIV-1. *Infect Dis Ther*. 2024;13(8):1877-1891. Available at: <https://pubmed.ncbi.nlm.nih.gov/38961048>.