

POSTPARTUM HIV MANAGEMENT AND FOLLOW-UP

Updated: March 31, 2026

Reviewed: March 31, 2026

Panel's Recommendations
<ul style="list-style-type: none"> Continuous antiretroviral therapy (ART) currently is recommended for all individuals with HIV to reduce the risk of disease progression and prevent the sexual transmission of HIV (AI). ART should be continued after giving birth (AI). Any plans for modifying ART after birth should be made in consultation with the individual and their HIV care provider, ideally before birth, taking into consideration the recommended regimens for nonpregnant adults (AII) and plans for future pregnancies. Because the immediate postpartum period poses unique challenges to ART adherence and retention in HIV care, arrangements for new or continued supportive services should be made throughout pregnancy and before postpartum hospital discharge (AII). Confirmatory testing should follow a positive HIV test during labor; see Pregnancy and Postpartum HIV Testing and Identification of Perinatal and Postnatal HIV Exposure. If testing confirms HIV infection, ART should be offered, and a supply of ART should be provided before postpartum hospital discharge to prevent treatment interruption (AII). Immediate linkage to HIV care and comprehensive follow-up also is needed (AII). When HIV is newly diagnosed in the intrapartum period, infants should begin presumptive HIV therapy, and ART should be provided before postpartum hospital discharge (AII) (see Antiretroviral Management of Infants With Antiretroviral Management of Infants With In Utero, Intrapartum, or Breastfeeding Exposure to HIV). In the context of HIV during pregnancy, evidence-based counseling should be provided to support shared decision-making about infant feeding options prior to and during pregnancy; counseling and plans for infant feeding should be reviewed again after birth (AIII) (see Preventing HIV Transmission During Infant Feeding). Clinicians should discuss future reproductive plans and timing, as well as the risks and benefits of conceiving while on specific antiretroviral medications (AII). The use of appropriate contraceptive options to prevent unintended pregnancy and optimal interpregnancy intervals should also be discussed (AII) (see Prepregnancy Counseling and Care). Contraceptive counseling should involve shared decision-making and should start during the prenatal period; a contraceptive plan should be developed before postpartum hospital discharge, as desired by the patient (AII).
<p>Rating of Recommendations: A = Strong; B = Moderate; C = Optional</p> <p>Rating of Evidence: I = One or more randomized trials with clinical outcomes and/or validated laboratory endpoints; II = One or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion</p>

The postpartum period provides an opportunity to review and optimize a patient's health care. Comprehensive medical care and supportive services are particularly important for patients with HIV and their families, who often face multiple medical and social challenges. Components of comprehensive care include the following services, as needed:

- Primary care, gynecologic/obstetric care, and HIV specialty care
- Subspecialty care as needed tailored to existing comorbidities

- Pediatric and pediatric HIV specialty care for the infant, including ongoing review of infant feeding plans and related prophylaxis and laboratory monitoring
- Sexual and reproductive health services, including contraception
- Mental health services
- Substance use prevention and treatment
- Supportive services
- Coordination of care through case management for the patient, their child (or children), and other family members
- Prevention of secondary transmission for partners with differing HIV status, including counseling on the use of condoms, antiretroviral therapy (ART) to maintain virologic suppression in the partner with HIV (i.e., treatment as prevention), and the potential use of pre-exposure prophylaxis (PrEP) by the partner without HIV (see [Pre-exposure Prophylaxis \[PrEP\] to Prevent HIV During Periconception, Antepartum, and Postpartum Periods](#))

Tailored supportive services can include screening for intimate partner violence; case management; child care; respite care; assistance with basic needs, such as housing, food, and transportation; peer counseling; and legal and advocacy services. Ideally, these services should begin before pregnancy and continue throughout pregnancy and the postpartum period.

Postpartum care should include immediate linkage to care, thorough medical assessment, counseling, and follow-up, with particular attention to new HIV diagnoses during labor or birth. This is important because intrapartum diagnosis of HIV has been negatively associated with HIV care outcomes in the postpartum period.^{1,2} The American College of Obstetricians and Gynecologists recommends contact with obstetrician-gynecologists within 3 weeks postpartum and that postpartum care be provided as an ongoing process based on individual needs rather than as a single postpartum visit.³ A follow-up appointment within 2 to 4 weeks of postpartum discharge is recommended with the health care provider managing HIV care, whether an obstetrician or an HIV specialist, particularly in the context of ART adherence challenges. Telemedicine visits should be considered when in-person appointment attendance is a concern.⁴

When care is not co-located or not within the same health care system, a case manager's role for care coordination becomes even more crucial. Case management improves the likelihood of achieving viral suppression and maintaining continuity of care.⁵ Alternative models of HIV care delivery—such as HIV-adapted group prenatal care—have been associated with higher retention in HIV care for women in the postpartum period.⁶ Ensuring continuity of ART between the antepartum and postpartum periods is especially critical. When HIV is newly diagnosed in the intrapartum period, infants should begin presumptive HIV therapy, and ART should be provided before postpartum hospital discharge. Special hospital programs may need to be established to support dispensing ART before discharge.

Patients taking testosterone or other medications and patients whose self-expression is less common may require additional support and linkage to care during the postpartum period.⁷⁻⁹

Postpartum Antiretroviral Therapy

ART should be continued postpartum. Decisions about changes to an ART regimen after birth should be made after discussion between the HIV care provider and patient, ideally before birth. Health care providers counseling on postpartum ART should factor in the possibility of future pregnancies (planned or unplanned) alongside the anticipated ART regimen, contraceptive choices,

and any unique drug–drug interactions in the postpartum period (see [Prenatal Counseling and Care](#) and [Appendix C: Antiretroviral Counseling Guide for Health Care Providers](#)).

ART is currently recommended for all individuals with HIV to reduce the risk of disease progression and to prevent transmission of HIV.¹⁰ The Strategic Timing of AntiRetroviral Treatment (START)¹¹ and TEMPRANO trials¹² were randomized clinical trials that demonstrated that early ART can reduce the risk of disease progression, even in individuals with CD4 T lymphocyte cell counts >500 cells/mm³, and the HIV Prevention Trials Network 052 randomized clinical trial demonstrated that effective ART can reduce the risk of sexual transmission of HIV to a partner without HIV by 100%.¹³ People with HIV who take ART as prescribed and achieve and maintain an undetectable viral load have effectively no risk of transmitting HIV through sex (i.e., Undetectable = Untransmittable).¹⁴

Postpartum counseling should emphasize the importance of lifelong ART. Several studies have demonstrated significant decreases in ART adherence postpartum.^{15–18} During the postpartum period, medical appointment follow-up may be difficult, including appointment adherence, which can affect ART adherence. Systematic monitoring of retention in HIV care is recommended for all individuals with HIV, but special attention is warranted during the postpartum period.

Adherence to Antiretroviral Therapy and Postpartum Depression

Symptoms of depression have been associated with lower ART adherence and viral suppression during pregnancy and the postpartum period.¹⁹ Furthermore, postpartum depression is common in the context of HIV.^{2,20–23} The U.S. Preventive Services Task Force recommends universal screening for postpartum depression²⁴ using a validated tool (e.g., the Edinburgh Postnatal Depression Scale); such screening is especially important when increased risk for postpartum depression and poor adherence is suspected. Enquiring about intimate partner violence, food or housing insecurity, and substance use is also important, as these factors have been associated with worsening depressive symptoms.²⁵ Counseling should emphasize that postpartum physical and psychological changes (and the stresses and demands of caring for a new baby) may make adherence more difficult and that additional support may be needed during this period.^{5,26–29} In addition, depression care with antidepressants and individual problem-solving therapy has shown to improve maternal functioning and infant care.²³

Poor adherence has been associated with virologic failure, the development of antiretroviral (ARV) drug resistance, and decreased long-term effectiveness of ART.^{30–32} If viral suppression is achieved by the time of birth, postpartum ART simplification to once-daily, co-formulated regimens or long-acting injectable ART—the usual preferred initial regimens for non-pregnant adults—could promote adherence during this challenging time. Efforts to maintain adequate adherence during the postpartum period may ensure effectiveness of therapy (see [Adherence to the Continuum of Care](#) in the [Adult and Adolescent Antiretroviral Guidelines](#)). When a protease inhibitor–based ART regimen is continued but at increased doses during pregnancy, available data suggest that doses can be reduced to standard doses immediately after birth. In addition, such behavioral interventions as peer support and text messaging may help improve retention in HIV care postpartum.³³

Secondary Sexual Transmission and Contraception

The postpartum period is a critical time for addressing safer sex practices to reduce secondary transmission of HIV to partners,³⁴ and clinicians should begin discussing these practices with the

patient during the prenatal period. Full, sustained viral suppression (undetectable viral load)—with or without reliable PrEP use by the partner—eliminates the risk of sexual transmission (see [Reproductive Options When One or Both Partners Have HIV](#) and [National Institutes of Health's Treatment as Prevention](#) guidance for more information). However, additional topics for discussion during counseling on prevention of secondary transmission to the partner without HIV should include condom use and the potential use of PrEP by the partner without HIV, both of which may play important roles in some clinical scenarios (e.g., where sustained ART adherence and viral load suppression may not be known with certainty, where patients desire additional forms of risk reduction).

Comprehensive pre-pregnancy counseling and contraception should be integrated into all health care visits, with special attention given to these topics during routine prenatal and postpartum visits. Lack of breastfeeding is associated with earlier return of fertility. Ovulation returns as early as 6 weeks postpartum, and it can occur earlier in some cases—even before resumption of menses—increasing the likelihood of pregnancy soon after birth.³⁵ HIV does not preclude the use of any contraceptive method (see [Table 3. Drug Interactions Between Antiretroviral Agents and Hormonal Contraceptives](#)). Contraceptive plans should not vary based on HIV status; however, because of the significantly higher prevalence of unplanned pregnancy in the context of HIV,³⁶ establishing contraceptive plans early, using shared decision-making, is important. If a long-acting reversible contraceptive (LARC)—such as an injectable, implant, or intrauterine device (IUD)—is desired, it should be inserted before postpartum hospital discharge or during the routine postpartum visit. If LARC insertion is planned for the postpartum visit and an interim contraceptive is desired, intramuscular depot medroxyprogesterone acetate is an option that can be given before postpartum hospital discharge. Counseling should advise against interpregnancy intervals shorter than 6 months; however, the contraceptive plan should be made using a shared decision-making approach.^{3,37,38}

The potential for drug–drug interactions between several ARV drugs and hormonal contraceptives is discussed in [Prenatal Counseling and Care](#) and [Table 3. Drug Interactions Between Antiretroviral Agents and Hormonal Contraceptives](#). A systematic review conducted for the World Health Organization summarized the research on hormonal contraception, IUD use, and risk of HIV infection and concluded that women with HIV can use all forms of contraception.^{39,40} This is consistent with Centers for Disease Control and Prevention recommendations advocating access to a broad range of effective contraceptive methods, including combined hormonal contraceptives, progestin-only pills, depot medroxyprogesterone acetate, and implants.⁴¹

Infant Feeding

In the context of HIV during pregnancy, evidence-based counseling should be provided to support shared decision-making about infant feeding options prior to and during pregnancy (see [Preventing HIV Transmission During Infant Feeding](#)). Counseling and plans for infant feeding should be reviewed again after birth.

Lactation Inhibition

In the absence of breastfeeding, symptoms related to breast engorgement can be very unpleasant in the days following labor and birth. Supportive measures—such as using acetaminophen or ibuprofen for pain control, alternating hot and cold compresses on the breasts, or wearing a tight-fitting bra—can help relieve symptoms related to breast engorgement.³ Although pharmacologic options for lactation inhibition generally are not used in the United States, recent data suggest cabergoline may be appropriate in some cases.^{42,43} Cabergoline is a dopamine agonist/ergot derivative that reduces

the production of prolactin; however, it is not approved by the U.S. Food and Drug Administration for lactation inhibition.^{44,45} Cabergoline is **contraindicated** for women with hypertension—including pregnancy-induced hypertension, preeclampsia, or eclampsia—or liver disease and for women being treated with antipsychotics or those who have a history of puerperal psychosis.⁴⁶ Bromocriptine, another dopamine agonist, is no longer used for lactation inhibition because of serious cardiovascular and neurologic complications associated with its use.⁴⁷

References

1. Momplaisir FM, Brady KA, Fekete T, et al. Time of HIV diagnosis and engagement in prenatal care impact virologic outcomes of pregnant women with HIV. *PLoS One*. 2015;10(7):e0132262. Available at: <https://pubmed.ncbi.nlm.nih.gov/26132142>.
2. Paioni P, Aebi-Popp K, Martinez de Tejada B, et al. Viral suppression and retention in HIV care during the postpartum period among women living with HIV: a longitudinal multicenter cohort study. *Lancet Reg Health Eur*. 2023;31:100656. Available at: <https://pubmed.ncbi.nlm.nih.gov/37303945>.
3. American College of Obstetricians and Gynecologists. ACOG committee opinion no. 736: optimizing postpartum care. *Obstet Gynecol*. 2018;131(5):e140-e150. Available at: <https://pubmed.ncbi.nlm.nih.gov/29683911>.
4. Whittington JR, Magann EF. Telemedicine in high-risk obstetrics. *Obstet Gynecol Clin North Am*. 2020;47(2):249-257. Available at: <https://pubmed.ncbi.nlm.nih.gov/32451016>.
5. Anderson EA, Momplaisir FM, Corson C, Brady KA. Assessing the impact of perinatal HIV case management on outcomes along the HIV care continuum for pregnant and postpartum women living with HIV, Philadelphia 2005–2013. *AIDS Behav*. 2017;21(9):2670-2681. Available at: <https://pubmed.ncbi.nlm.nih.gov/28176167>.
6. McKinney J, Jackson J, Sangi-Haghpeykar H, et al. HIV-adapted group prenatal care: assessing viral suppression and postpartum retention in care. *AIDS Patient Care STDS*. 2021;35(2):39-46. Available at: <https://pubmed.ncbi.nlm.nih.gov/33571047>.
7. American College of Obstetricians Gynecologists 'Committee on Gynecologic Practice, American College of Obstetricians Gynecologists 'Committee on Health Care for Underserved Women. Health care for transgender and gender diverse individuals: ACOG committee opinion, number 823. *Obstet Gynecol*. 2021;137(3):e75-e88. Available at: <https://pubmed.ncbi.nlm.nih.gov/33595253>.
8. MacDonald T, Noel-Weiss J, West D, et al. Transmasculine individuals 'experiences with lactation, chestfeeding, and gender identity: a qualitative study. *BMC Pregnancy Childbirth*. 2016;16:106. Available at: <https://pubmed.ncbi.nlm.nih.gov/27183978>.
9. MacLean LR. Preconception, pregnancy, birthing, and lactation needs of transgender men. *Nurs Womens Health*. 2021;25(2):129-138. Available at: <https://pubmed.ncbi.nlm.nih.gov/33651985>.
10. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in adults and adolescents with HIV. 2025. Available at: <https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-arv/whats-new>.
11. INSIGHT START Study Group, Lundgren JD, Babiker AG, et al. Initiation of antiretroviral therapy in early asymptomatic HIV infection. *N Engl J Med*. 2015;373(9):795-807. Available at: <https://pubmed.ncbi.nlm.nih.gov/26192873>.
12. TEMPRANO ANRS Study Group, Daniel C, Moh R, et al. A trial of early antiretrovirals and isoniazid preventive therapy in Africa. *N Engl J Med*. 2015;373(9):808-822. Available at: <https://pubmed.ncbi.nlm.nih.gov/26193126>.

13. Cohen MS, Chen YQ, McCauley M, et al. Antiretroviral therapy for the prevention of HIV-1 transmission. *N Engl J Med*. 2016;375(9):830-839. Available at: <https://pubmed.ncbi.nlm.nih.gov/27424812>.
14. Centers for Disease Control and Prevention. HIV transmission. 2019. Available at: <https://www.cdc.gov/hiv/basics/transmission.html>.
15. Kreitchmann R, Coelho DF, Kakehasi FM, et al. Long-term postpartum adherence to antiretroviral drugs among women in Latin America. *Int J STD AIDS*. 2016;27(5):377-386. Available at: <https://pubmed.ncbi.nlm.nih.gov/25931238>.
16. Adams JW, Brady KA, Michael YL, et al. Postpartum engagement in HIV care: an important predictor of long-term retention in care and viral suppression. *Clin Infect Dis*. 2015;61(12):1880-1887. Available at: <https://pubmed.ncbi.nlm.nih.gov/26265499>.
17. Nachega JB, Uthman OA, Anderson J, et al. Adherence to antiretroviral therapy during and after pregnancy in low-income, middle-income, and high-income countries: a systematic review and meta-analysis. *AIDS*. 2012;26(16):2039-2052. Available at: <https://pubmed.ncbi.nlm.nih.gov/22951634>.
18. Ojukwu EN, Cianelli R, Rodriguez NV, et al. Predictors and social determinants of HIV treatment engagement among post-partum Black women living with HIV in southeastern United States. *J Adv Nurs*. 2023;79(11):4365-4380. Available at: <https://pubmed.ncbi.nlm.nih.gov/37243385>.
19. Momplaisir F, Hussein M, Kacanek D, et al. Perinatal depressive symptoms, HIV suppression, and the underlying role of ART adherence: a longitudinal mediation analysis in the IMPAACT P1025 cohort. *Clin Infect Dis*. 2021;73(8):1379-1387. Available at: <https://pubmed.ncbi.nlm.nih.gov/33982083>.
20. Ion A, Wagner AC, Greene S, et al. HIV-related stigma in pregnancy and early postpartum of mothers living with HIV in Ontario, Canada. *AIDS Care*. 2017;29(2):137-144. Available at: <https://pubmed.ncbi.nlm.nih.gov/27449254>.
21. Wielding S, Scott A. What women want: social characteristics, gender-based violence and social support preferences in a cohort of women living with HIV. *Int J STD AIDS*. 2017;28(5):486-490. Available at: <https://pubmed.ncbi.nlm.nih.gov/27270691>.
22. Gauthreaux C, Negron J, Castellanos D, et al. The association between pregnancy intendedness and experiencing symptoms of postpartum depression among new mothers in the United States, 2009 to 2011: a secondary analysis of PRAMS data. *Medicine (Baltimore)*. 2017;96(6):e5851. Available at: <https://pubmed.ncbi.nlm.nih.gov/28178128>.
23. Wagner GJ, Ghosh-Dastidar B, Faherty L, et al. Effects of M-DEPTH model of depression care on maternal functioning and infant developmental outcomes in the six months post delivery: results from a cluster randomized controlled trial. *Womens Reprod Health (Phila)*. 2024;11(2):313-328. Available at: <https://pubmed.ncbi.nlm.nih.gov/39081835>.
24. O'Connor E, Rossom RC, Henninger M, et al. Primary care screening for and treatment of depression in pregnant and postpartum women: evidence report and systematic review for the US Preventive Services Task Force. *JAMA*. 2016;315(4):388-406. Available at: <https://pubmed.ncbi.nlm.nih.gov/26813212>.

25. Larsen AM, Osborn L, Ronen K, et al. Trajectories of depression symptoms from pregnancy through 24 months postpartum among Kenyan women living with HIV. *J Acquir Immune Defic Syndr*. 2022;90(5):473-481. Available at: <https://pubmed.ncbi.nlm.nih.gov/35394987>.
26. Cohn SE, Umbleja T, Mrus J, et al. Prior illicit drug use and missed prenatal vitamins predict nonadherence to antiretroviral therapy in pregnancy: adherence analysis A5084. *AIDS Patient Care STDS*. 2008;22(1):29-40. Available at: <https://pubmed.ncbi.nlm.nih.gov/18442305>.
27. Ickovics JR, Wilson TE, Royce RA, et al. Prenatal and postpartum zidovudine adherence among pregnant women with HIV: results of a MEMS substudy from the perinatal guidelines evaluation project. *J Acquir Immune Defic Syndr*. 2002;30(3):311-315. Available at: <https://pubmed.ncbi.nlm.nih.gov/12131568>.
28. Bardeguet AD, Lindsey JC, Shannon M, et al. Adherence to antiretrovirals among US women during and after pregnancy. *J Acquir Immune Defic Syndr*. 2008;48(4):408-417. Available at: <https://pubmed.ncbi.nlm.nih.gov/18614923>.
29. Buchberg MK, Fletcher FE, Vidrine DJ, et al. A mixed-methods approach to understanding barriers to postpartum retention in care among low-income, HIV-infected women. *AIDS Patient Care STDS*. 2015;29(3):126-132. Available at: <https://pubmed.ncbi.nlm.nih.gov/25612217>.
30. Paterson DL, Swindells S, Mohr J, et al. Adherence to protease inhibitor therapy and outcomes in patients with HIV infection. *Ann Intern Med*. 2000;133(1):21-30. Available at: <https://pubmed.ncbi.nlm.nih.gov/10877736>.
31. Le Moing V, Chêne G, Carrieri MP, et al. Clinical, biologic, and behavioral predictors of early immunologic and virologic response in HIV-infected patients initiating protease inhibitors. *J Acquir Immune Defic Syndr*. 2001;27(4):372-376. Available at: <https://pubmed.ncbi.nlm.nih.gov/11468425>.
32. Murri R, Ammassari A, Gallicano K, et al. Patient-reported nonadherence to HAART is related to protease inhibitor levels. *J Acquir Immune Defic Syndr*. 2000;24(2):123-128. Available at: <https://pubmed.ncbi.nlm.nih.gov/10935687>.
33. Abuogi LL, Onono M, Odeny TA, et al. Effects of behavioural interventions on postpartum retention and adherence among women with HIV on lifelong ART: the results of a cluster randomized trial in Kenya (the MOTIVATE trial). *J Int AIDS Soc*. 2022;25(1):e25852. Available at: <https://pubmed.ncbi.nlm.nih.gov/35041776>.
34. Cates W Jr, Steiner MJ. Dual protection against unintended pregnancy and sexually transmitted infections: what is the best contraceptive approach? *Sex Transm Dis*. 2002;29(3):168-174. Available at: <https://pubmed.ncbi.nlm.nih.gov/11875378>.
35. Jackson E, Glasier A. Return of ovulation and menses in postpartum nonlactating women: a systematic review. *Obstet Gynecol*. 2011;117(3):657-662. Available at: <https://pubmed.ncbi.nlm.nih.gov/21343770>.
36. Sutton MY, Patel R, Frazier EL. Unplanned pregnancies among HIV-infected women in care-United States. *J Acquir Immune Defic Syndr*. 2014;65(3):350-358. Available at: <https://pubmed.ncbi.nlm.nih.gov/24189153>.

37. Sholapurkar SL. Is there an ideal interpregnancy interval after a live birth, miscarriage or other adverse pregnancy outcomes? *J Obstet Gynaecol*. 2010;30(2):107-110. Available at: <https://pubmed.ncbi.nlm.nih.gov/20143964>.
38. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. *JAMA*. 2006;295(15):1809-1823. Available at: <https://pubmed.ncbi.nlm.nih.gov/16622143>.
39. World Health Organization. Review of priorities in research: hormonal contraception and IUDs and HIV infection. 2010. Geneva, Switzerland. Available at: <https://iris.who.int/server/api/core/bitstreams/17233811-f9e6-4974-9ecc-73f3ea348f8d/content>.
40. Polis CB, Curtis KM. Use of hormonal contraceptives and HIV acquisition in women: a systematic review of the epidemiological evidence. *Lancet Infect Dis*. 2013;13(9):797-808. Available at: <https://pubmed.ncbi.nlm.nih.gov/23871397>.
41. Curtis KM, Tepper NK, Jatlaoui TC, et al. U.S. medical eligibility criteria for contraceptive use, 2016. *MMWR Recomm Rep*. 2016;65(3):1-103. Available at: <https://pubmed.ncbi.nlm.nih.gov/27467196>.
42. Tulloch KJ, Dodin P, Tremblay-Racine F, et al. Cabergoline: a review of its use in the inhibition of lactation for women living with HIV. *J Int AIDS Soc*. 2019;22(6):e25322. Available at: <https://pubmed.ncbi.nlm.nih.gov/31183987>.
43. Harris K, Murphy KE, Horn D, et al. Safety of cabergoline for postpartum lactation inhibition or suppression: a systematic review. *J Obstet Gynaecol Can*. 2020;42(3):308-315 e320. Available at: <https://pubmed.ncbi.nlm.nih.gov/31285168>.
44. Henkel A, Johnson SA, Reeves MF, et al. Cabergoline for lactation inhibition after second-trimester abortion or pregnancy loss: a randomized controlled trial. *Obstet Gynecol*. 2023;141(6):1115-1123. Available at: <https://pubmed.ncbi.nlm.nih.gov/37486652>.
45. National Institute of Child Health and Human Development. Cabergoline. *Drugs and Lactation Database (LactMed®)*. 2023. Available at: <https://pubmed.ncbi.nlm.nih.gov/30000386>.
46. Dostinex (cabergoline) [package insert]. Food and Drug Administration. 2019. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2011/020664s011lbl.pdf.
47. Bromocriptine mesylate (Parlodel) for the prevention of physiological lactation; opportunity for a hearing on a proposal to withdraw approval of the indication [package insert]. Food and Drug Administration. 1994. Available at: <https://www.govinfo.gov/content/pkg/FR-1994-08-23/html/94-20562.htm>.