

Pregnant Women Living with HIV Who Have Never Received Antiretroviral Drugs (Antiretroviral Naive) (Last updated December 12, 2019; last reviewed December 12, 2019)

Panel's Recommendations

- Antiretroviral therapy (ART) is recommended for all pregnant women living with HIV to reduce the risk of perinatal transmission of HIV and to optimize the health of the mother (AI). Initiating ART as soon as possible in pregnant women who have never received antiretroviral (ARV) drugs is recommended, based on data demonstrating that earlier virologic suppression is associated with a lower risk of transmission (AII).
- The results of HIV drug-resistance studies should guide the selection of antiretroviral (ARV) regimens in women whose HIV RNA levels are above the threshold for resistance testing (i.e., >500 copies/mL to 1,000 copies/mL), unless drug-resistance studies have already been performed (see [Antiretroviral Drug Resistance and Resistance Testing in Pregnancy](#)) (AII). When ART is initiated before the results of the drug-resistance assays are available, the ARV regimen should be modified, if necessary, based on the resistance assay results (BIII).
- ARV regimens that are Preferred for the treatment of pregnant women with HIV who are ARV-naive include: a dual-nucleoside reverse transcriptase inhibitor combination (abacavir plus lamivudine or tenofovir disoproxil fumarate plus either emtricitabine or lamivudine) and either a ritonavir-boosted protease inhibitor (atazanavir/ritonavir or darunavir/ritonavir) or an integrase strand transfer inhibitor (dolutegravir [irrespective of trimester] or raltegravir; see [Table 4](#) and [Updated Guidance about the Use of Dolutegravir in Pregnancy Recommendations for the Use of Antiretroviral Drugs During Pregnancy](#)) (AIII).
- The Panel on Treatment of Pregnant Women with HIV Infection and Prevention of Perinatal Transmission emphasizes the importance of counseling and informed decision-making with regard to all ARV regimens for people living with HIV (AIII). See [Appendix D: Dolutegravir Counseling Guide for Health Care Providers](#) for more information.

Rating of Recommendations: A = Strong; B = Moderate; C = Optional

Rating of Evidence: I = One or more randomized trials with clinical outcomes and/or validated laboratory endpoints; II = One or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion

Pregnant women living with HIV should receive standard clinical, immunologic, and virologic evaluations. Consistent with the principles of HIV treatment for nonpregnant adults, clinicians should discuss treatment options with pregnant women and offer antiretroviral (ARV) regimens that contain at least three drugs. These regimens reduce the risk of perinatal HIV transmission and optimize the woman's health. Use of an ARV regimen that successfully reduces plasma HIV RNA to undetectable levels substantially lowers the risk of perinatal transmission of HIV, minimizes the need to consider elective cesarean delivery as an intervention to reduce the risk of transmission, and reduces the risk of ARV drug resistance in the mother.

Decisions about the timing and management of antiretroviral therapy (ART) in women who have not previously received ART should be guided by several key principles:

A suppressed viral load at the time of delivery markedly reduces perinatal transmission risk.

In an analysis of 12,486 infants delivered by women with HIV between 2000 and 2011 in the United Kingdom and Ireland, the overall perinatal transmission rate declined from 2.1% in 2000 and 2001 to 0.46% in 2010 and 2011. The transmission risk was significantly lower in women with viral loads <50 copies/mL (0.09%) than in women with viral loads of 50 copies/mL to 399 copies/mL (1.0%), regardless of the type of ARV regimen used or the mode of infant delivery.¹ The decline in perinatal transmission rates was attributed to the increasing number of women on ART at the time of conception and reductions in the proportion of women who either initiated ART late in pregnancy or who never received ART prior to delivery.

Initiating ART early increases the likelihood that a woman will achieve viral suppression by the time of delivery, further reducing transmission risk.

Although most perinatal transmission events occur late in pregnancy or during delivery, recent analyses suggest that early control of viral replication may be important in preventing transmission. In the prospective multicenter French Perinatal Cohort, both maternal viral load at delivery and the timing of ART initiation

were independently associated with perinatal transmission rate. For women who achieved viral loads <50 copies/mL at the time of delivery, transmission risks were 0.9% with third-trimester ART initiation, 0.5% with second-trimester initiation, 0.2% with first-trimester initiation, and 0% (of more than 2,500 infants) with preconception ART initiation. Regardless of when ART was initiated, the perinatal transmission rate was higher for women with viral loads of 50 copies/mL to 400 copies/mL near delivery than for those with <50 copies/mL, and higher still for women with viral loads >400 copies/mL at delivery (4.4% for women who initiated ART in the third trimester and who had viral loads >400 copies/mL at delivery).²

In an earlier publication that reported on the same cohort, lack of early and sustained control of maternal viral load appeared to be strongly associated with residual perinatal transmission of HIV.³ Similar data from Canada in 1,707 pregnant women with HIV who were followed between 1997 and 2010 showed that the risk of perinatal transmission was 1% in all mothers who received ART and 0.4% if ART was taken for more than 4 weeks.⁴

These data suggest that ART should be initiated **as early as possible** in ARV-naive women, because early and sustained control of HIV viral replication is associated with a decreased risk of transmission. Other studies have demonstrated that baseline viral load is significantly associated with the likelihood of viral suppression by delivery; thus, prompt initiation of ART is particularly important in pregnant women who have high baseline viral loads.^{5-7,8}

The benefits of initiating ART early in pregnancy generally outweigh the risks.

The susceptibility of fetuses to the potential adverse effects of drugs is dependent on multiple factors, including the gestational age of the fetus at **the time of medication** exposure (see [Teratogenicity](#) and [Combination Antiretroviral Drug Regimens and Maternal and Neonatal Outcomes](#)). The effects of taking ARV drugs during pregnancy are not fully known; however, in general, the data from observational studies on the incidence of birth defects among fetuses/infants of women who received ARV regimens during pregnancy have been reassuring. There have been no differences between the rates of birth defects among infants with first-trimester exposures to most ARV drugs and the rates among infants with later gestational exposures or the rates reported in the general population.⁹⁻¹² **Please see [Teratogenicity](#) for a more detailed discussion of the adverse events that are associated with the use of specific ARV drugs, including dolutegravir (DTG).** The decision about when to initiate ART should be discussed by health care providers and their patients. The discussion should include an assessment of a woman's health status and the risks and benefits to her health and the potential risks and benefits to the fetus.

ARV drugs further reduce transmission risk through infant pre-exposure and post-exposure prophylaxis.

Although rates of perinatal transmission are low in women with undetectable or low HIV RNA levels, there is no threshold below which lack of transmission can be ensured.¹³⁻¹⁵ ARV drugs reduce the risk of perinatal transmission of HIV through a number of different mechanisms. Although lowering maternal antenatal viral load is an important component of preventing transmission in women with higher viral loads, maternal ART use reduces transmission even in women with low viral loads.¹⁶⁻²⁰ Additional mechanisms that reduce the risk of perinatal HIV transmission include pre-exposure prophylaxis (PrEP) and post-exposure prophylaxis (PEP) for the infant.

With PrEP, the passage of an ARV drug across the placenta produces drug levels that inhibit viral replication in the fetus, particularly during the birth process when there is intensive viral exposure. Therefore, whenever possible, ARV regimens initiated during pregnancy should include a nucleoside reverse transcriptase inhibitor (NRTI) with high transplacental passage, such as lamivudine (3TC), emtricitabine (FTC), tenofovir disoproxil fumarate (TDF), or abacavir (ABC) (see [Table 8](#)).²¹⁻²⁴ With PEP, ARV drugs are administered to the infant after birth (see [Antiretroviral Management of Newborns with Perinatal HIV Exposure or HIV Infection](#)).

Specific ARV regimens are *Preferred* for use in pregnancy.

Tables [4](#) and [5](#) outline the ARV regimens that are designated by the Panel on Treatment of Pregnant Women with HIV Infection and Prevention of Perinatal Transmission as *Preferred* for treatment of pregnant women with HIV who have never received ARV drugs, as well as for women who are continuing or restarting ART

in pregnancy or women who are trying to conceive. Drugs or drug combinations are designated as *Preferred* for therapy in pregnant women when clinical trial data in adults have demonstrated efficacy and durability with acceptable toxicity and ease of use, and pregnancy-specific pharmacokinetic (PK) data are available to guide dosing. In addition, **the available data must suggest a favorable risk-benefit balance for the drug or drug combination compared to other ARV drug options; the assessment of risks and benefits should incorporate outcomes for women, fetuses, and infants. Some Preferred drugs or regimens may have minimal toxicity or teratogenicity risks that are offset by other advantages for women with HIV who are pregnant or who are trying to conceive.** Therefore, it is important to read all the information on each drug in the Perinatal Guidelines before administering any of these medications to patients (also see [Appendix B: Supplement: Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy](#)). *Preferred* regimens include a dual-NRTI combination (ABC plus 3TC or TDF plus FTC or 3TC) used with either a ritonavir-boosted protease inhibitor (PI; atazanavir/ritonavir or darunavir/ritonavir) or an integrase strand transfer inhibitor (INSTI; DTG or raltegravir [RAL]).

DTG is considered a *Preferred* INSTI for ART-naïve women, **irrespective of trimester.** It is a recommended option for an initial ARV regimen in nonpregnant adults, and there are sufficient data about the efficacy and safety of DTG when this drug is initiated during pregnancy.²⁵⁻²⁸ **Maternal use of DTG at the time of conception or in early pregnancy has been associated with an increase in the risk of neural tube defects (NTDs) in infants. However, this risk is small, and the decision about which ARV drugs to use during pregnancy should be made by a woman after discussing the known and potential benefits and risks to her and her fetus (infant).** DTG is an *Alternative* medication for women who are trying to conceive, due to concerns about NTDs that were observed in infants born to women who conceived while receiving DTG (see [Updated Guidance about the Use of Dolutegravir in Pregnancy](#) in [Recommendations for Use of Antiretroviral Drugs During Pregnancy](#)).

RAL is also a *Preferred* INSTI for ARV-naïve women, and the amount of efficacy and safety data for RAL in pregnant women is increasing. The selection of drugs for an ARV regimen should be based on individual patient characteristics and needs (see [Table 4](#)).

RAL or DTG have been suggested for use when ART is initiated late in pregnancy, particularly for women who have high viral loads, because of their ability to rapidly suppress viral load (a decrease of approximately 2 log₁₀ copies/mL occurs by Week 2 of therapy with these drugs).²⁹⁻³³ **In two open-label, randomized clinical trials in women who presented for treatment late in pregnancy, viral decline was more rapid and a greater proportion of women reached viral suppression targets when using INSTI-based regimens with RAL (IMPAACT 1081) or DTG (DolPHIN-2 study) than efavirenz-based ART.**^{28,34} DTG is *Preferred* for treatment of acute infection during pregnancy irrespective of trimester because it has a higher barrier to resistance than RAL and can be administered once daily. Because RAL has a lower barrier to resistance than DTG, it **is not recommended** for use during acute infection, when viral loads are expected to be high (see [Acute HIV Infection](#)). For a discussion regarding the addition of DTG or RAL to current ARV regimens, see [Lack of Viral Suppression](#).

Resistance tests should be performed, but ART initiation should not be delayed while waiting for results.

Standard ARV drug-resistance testing should be performed before starting an ARV regimen when plasma HIV RNA levels are above the threshold for resistance testing (i.e., >500 copies/mL to 1,000 copies/mL). INSTI-resistance testing is not routinely recommended, but it should be performed for women who are at risk for INSTI resistance (e.g., women with partners who were treated with INSTIs or women who had prior treatment that included INSTIs; see [Antiretroviral Drug Resistance and Resistance Testing in Pregnancy](#)). For details regarding genotypic and phenotypic resistance testing, see the [Adult and Adolescent Antiretroviral Guidelines](#). Given the association between earlier viral suppression and lower risk of perinatal transmission, ART should be initiated as soon as possible in pregnant women who have never received ARV drugs, without waiting for the results of resistance testing. The regimen can be modified, if required, when test results return. Either a PI-based or an INSTI-based ARV regimen can be considered when the results of resistance testing are not available to inform the selection of ARV drugs, because clinically significant resistance to PIs and INSTIs is uncommon in ARV-naïve individuals.

Regimens other than combination (three-drug) ART are not recommended.

The use of zidovudine (ZDV) monotherapy during pregnancy **is no longer recommended**, because ART provides clear health benefits to the mother and helps prevent perinatal HIV transmission. In the past, the use of ZDV monotherapy during pregnancy for prophylaxis of perinatal transmission was an option for women who had low viral loads (i.e., <1,000 copies/mL) on no ARV drugs. Although the Adult and Adolescent Antiretroviral Guidelines recommend some two-drug ARV regimens in certain clinical circumstances, two-drug ARV regimens **are not recommended** for use in pregnant women.

All pregnant women with HIV should be counseled that the use of ART is recommended, regardless of viral load, to optimally reduce the risk of perinatal transmission. If, after counseling, a woman chooses to forgo the use of ARV drugs during pregnancy, this decision should be re-addressed during subsequent medical appointments. The [Perinatal HIV Hotline](http://www.hivhotline.org) (1-800-439-4079) can provide information to assist with the discussion.

ARV regimens can be modified postpartum.

ARV regimens that were initiated during pregnancy can be modified after delivery. Women may be able to use some simplified regimens that could not be used during pregnancy because the pregnancy, safety, and/or PK data for those regimens were insufficient. Decisions regarding whether to continue an ARV regimen or which specific ARV agents to use postpartum should be made by women after they have discussed their options with their HIV care providers. These decisions should take several factors into consideration, including the current adult ART recommendations, a woman's plans for contraceptive use and future pregnancies, and individual adherence considerations and medication preferences (see [General Principles Regarding Use of Antiretroviral Drugs during Pregnancy](#)).

References

1. Townsend CL, Byrne L, Cortina-Borja M, et al. Earlier initiation of ART and further decline in mother-to-child HIV transmission rates, 2000-2011. *AIDS*. 2014;28(7):1049-1057. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24566097>.
2. Mandelbrot L, Tubiana R, Le Chenadec J, et al. No perinatal HIV-1 transmission from women with effective antiretroviral therapy starting before conception. *Clin Infect Dis*. 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26197844>.
3. Tubiana R, Le Chenadec J, Rouzioux C, et al. Factors associated with mother-to-child transmission of HIV-1 despite a maternal viral load <500 copies/ml at delivery: a case-control study nested in the French perinatal cohort (EPF-ANRS CO1). *Clin Infect Dis*. 2010;50(4):585-596. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20070234>.
4. Forbes JC, Alimenti AM, Singer J, et al. A national review of vertical HIV transmission. *AIDS*. 2012;26(6):757-763. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22210635>.
5. Read PJ, Mandalia S, Khan P, et al. When should HAART be initiated in pregnancy to achieve an undetectable HIV viral load by delivery? *AIDS*. 2012;26(9):1095-1103. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22441248>.
6. Katz IT, Shapiro R, Li D, et al. Risk factors for detectable HIV-1 RNA at delivery among women receiving highly active antiretroviral therapy in the women and infants transmission study. *J Acquir Immune Defic Syndr*. 2010;54(1):27-34. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20065861>.
7. Aziz N, Sokoloff A, Kornak J, et al. Time to viral load suppression in antiretroviral-naive and -experienced HIV-infected pregnant women on highly active antiretroviral therapy: implications for pregnant women presenting late in gestation. *BJOG*. 2013. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23924192>.
8. Myer L, Phillips TK, McIntyre JA, et al. HIV viraemia and mother-to-child transmission risk after antiretroviral therapy initiation in pregnancy in Cape Town, South Africa. *HIV Med*. 2017;18(2):80-88. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27353189>.
9. da Costa TP, Machado ES, et al. Malformations among HIV vertically exposed newborns—results from a Brazilian cohort study. Presented at: 6th IAS Conference on HIV Pathogenesis and Treatment and Prevention. 2011. Rome, Italy.
10. Watts DH, Huang S, Culnane M, et al. Birth defects among a cohort of infants born to HIV-infected women on antiretroviral medication. *J Perinat Med*. 2011;39(2):163-170. Available at: <http://www.ncbi.nlm.nih.gov/>

pubmed/21142844.

11. Knapp KM, Brogly SB, Muenz DG, et al. Prevalence of congenital anomalies in infants with *in utero* exposure to antiretrovirals. *Pediatr Infect Dis J*. 2012;31(2):164-170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21983213>.
12. Floridia M, Mastroiacovo P, Tamburrini E, et al. Birth defects in a national cohort of pregnant women with HIV infection in Italy, 2001-2011. *BJOG*. 2013;120(12):1466-1475. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23721372>.
13. Cooper ER, Charurat M, Mofenson L, et al. Combination antiretroviral strategies for the treatment of pregnant HIV-1-infected women and prevention of perinatal HIV-1 transmission. *J Acquir Immune Defic Syndr*. 2002;29(5):484-494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11981365>.
14. Mofenson LM, Lambert JS, Stiehm ER, et al. Risk factors for perinatal transmission of human immunodeficiency virus type 1 in women treated with zidovudine. Pediatric AIDS Clinical Trials Group Study 185 Team. *N Engl J Med*. 1999;341(6):385-393. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10432323>.
15. Garcia PM, Kalish LA, Pitt J, et al. Maternal levels of plasma human immunodeficiency virus type 1 RNA and the risk of perinatal transmission. Women and Infants Transmission Study Group. *N Engl J Med*. 1999;341(6):394-402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10432324>.
16. Ioannidis JP, Abrams EJ, Ammann A, et al. Perinatal transmission of human immunodeficiency virus type 1 by pregnant women with RNA virus loads <1000 copies/ml. *J Infect Dis*. 2001;183(4):539-545. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11170978>.
17. Wade NA, Birkhead GS, Warren BL, et al. Abbreviated regimens of zidovudine prophylaxis and perinatal transmission of the human immunodeficiency virus. *N Engl J Med*. 1998;339(20):1409-1414. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9811915>.
18. Jackson JB, Musoke P, Fleming T, et al. Intrapartum and neonatal single-dose nevirapine compared with zidovudine for prevention of mother-to-child transmission of HIV-1 in Kampala, Uganda: 18-month follow-up of the HIVNET 012 randomised trial. *Lancet*. 2003;362(9387):859-868. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/13678973>.
19. Petra Study Team. Efficacy of three short-course regimens of zidovudine and lamivudine in preventing early and late transmission of HIV-1 from mother to child in Tanzania, South Africa, and Uganda (Petra study): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2002;359(9313):1178-1186. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11955535>.
20. Moodley D, Moodley J, Coovadia H, et al. A multicenter randomized controlled trial of nevirapine versus a combination of zidovudine and lamivudine to reduce intrapartum and early postpartum mother-to-child transmission of human immunodeficiency virus type 1. *J Infect Dis*. 2003;187(5):725-735. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12599045>.
21. Hirt D, Urien S, Rey E, et al. Population pharmacokinetics of emtricitabine in human immunodeficiency virus type 1-infected pregnant women and their neonates. *Antimicrob Agents Chemother*. 2009;53(3):1067-1073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19104016>.
22. Hirt D, Urien S, Ekouevi DK, et al. Population pharmacokinetics of tenofovir in HIV-1-infected pregnant women and their neonates (ANRS 12109). *Clin Pharmacol Ther*. 2009;85(2):182-189. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18987623>.
23. Moodley D, Pillay K, Naidoo K, et al. Pharmacokinetics of zidovudine and lamivudine in neonates following coadministration of oral doses every 12 hours. *J Clin Pharmacol*. 2001;41(7):732-741. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11452705>.
24. Wade NA, Unadkat JD, Huang S, et al. Pharmacokinetics and safety of stavudine in HIV-infected pregnant women and their infants: pediatric AIDS clinical trials group protocol 332. *J Infect Dis*. 2004;190(12):2167-2174. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15551216>.
25. Zash R, Jacobson DL, Diseko M, et al. Comparative safety of dolutegravir-based or efavirenz-based antiretroviral treatment started during pregnancy in Botswana: an observational study. *Lancet Glob Health*. 2018;6(7):e804-e810. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29880310>.
26. Zash R, Holmes L, Makhema J, et al. Surveillance for neural tube defects following antiretroviral exposure from conception. Presented at: 22nd International AIDS Conference. 2018. Amsterdam, Netherlands. Available at: <http://www>.

natap.org/2018/IAC/IAC_52.htm.

27. Zash R, Holmes L, Diseko M, et al. Neural-tube defects and antiretroviral treatment regimens in Botswana. *N Engl J Med*. 2019;381(9):827-840. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31329379>.
28. Kintu K, Malaba T, Nakibuka J, et al. Rct of dolutegravir vs efavirenz-based therapy initiated in late pregnancy: DolPHIN-2. Abstract 40. Presented at: Conference on Retroviruses and Opportunistic Infections. 2019. Seattle, Washington. Available at: <http://www.croiconference.org/sessions/rct-dolutegravir-vs-efavirenz-based-therapy-initiated-late-pregnancy-dolphin-2>.
29. Grinsztejn B, Nguyen BY, Katlama C, et al. Safety and efficacy of the HIV-1 integrase inhibitor raltegravir (MK-0518) in treatment-experienced patients with multidrug-resistant virus: a Phase II randomised controlled trial. *Lancet*. 2007;369(9569):1261-1269. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17434401>.
30. Papendorp SG, van den Berk GE. Preoperative use of raltegravir-containing regimen as induction therapy: very rapid decline of HIV-1 viral load. *AIDS*. 2009;23(6):739. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19279447>.
31. Pinnetti C, Baroncelli S, Villani P, et al. Rapid HIV-RNA decline following addition of raltegravir and tenofovir to ongoing highly active antiretroviral therapy in a woman presenting with high-level HIV viraemia at week 38 of pregnancy. *J Antimicrob Chemother*. 2010;65(9):2050-2052. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20630894>.
32. McKeown DA, Rosenvinge M, Donaghy S, et al. High neonatal concentrations of raltegravir following transplacental transfer in HIV-1 positive pregnant women. *AIDS*. 2010;24(15):2416-2418. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20827058>.
33. Orrell C, Kintu JA, et al. DolPHIN-1: Randomised controlled trial of dolutegravir(DTG)-versus efavirenz(EFV)-based therapy in mothers initiating antiretroviral treatment in late pregnancy. Presented at: 22nd International AIDS Conference. 2018. Amsterdam, Netherlands. Available at: http://www.natap.org/2018/IAC/IAC_30.htm.
34. Mirochnick M, Shapiro D, Morrison L, et al. Randomized trial of raltegravir-art vs efavirenz-art when initiated during pregnancy. Abstract 39. Presented at: Conference on Retroviruses and Opportunistic Infections. 2019. Seattle, Washington. Available at: <http://www.croiconference.org/sessions/randomized-trial-raltegravir-art-vs-efavirenz-art-when-initiated-during-pregnancy>.