

Appendix B, Table 1. Coformulated Single-Tablet Regimens (Last updated July 10, 2019; last reviewed December 18, 2019)

The following table includes dose recommendations for FDA-approved STR products. Please see the class-specific drug characteristics tables ([Appendix B, Tables 3 to 6](#)) for details about the individual drugs included in these STR products, including information on elimination and metabolic pathways, serum and intracellular half-lives, and adverse effects. The STR products in this table are listed by drug class and arranged in **alphabetical order** by trade name within each class.

Trade Name (Abbreviations)	ARV Drugs Included in the STR	Dosing Recommendation ^a
INSTI plus Two NRTIs		
Biktarvy (BIC/TAF/FTC)	Bictegravir 50 mg/tenofovir alafenamide 25 mg/emtricitabine 200 mg	One tablet once daily
Genvoya (EVG/c/TAF/FTC)	Elvitegravir 150 mg/cobicistat 150 mg/tenofovir alafenamide 10 mg/emtricitabine 200 mg	One tablet once daily with food
Stribild (EVG/c/TDF/FTC)	Elvitegravir 150 mg/cobicistat 150 mg/tenofovir disoproxil fumarate 300 mg/emtricitabine 200 mg	One tablet once daily with food
Triumeq (DTG/ABC/3TC)	Dolutegravir 50 mg/abacavir 600 mg/lamivudine 300 mg	One tablet once daily
INSTI plus One NRTI		
Dovato (DTG/3TC)	Dolutegravir 50 mg/lamivudine 300 mg	One tablet once daily
PI plus Two NRTIs		
Symtuza (DRV/c/TAF/FTC)	Darunavir 800 mg/cobicistat 150 mg/tenofovir alafenamide 10 mg/emtricitabine 200 mg	One tablet once daily with food
NNRTI plus Two NRTIs		
Atripla (EFV/TDF/FTC)	Efavirenz 600 mg/tenofovir disoproxil fumarate 300 mg/emtricitabine 200 mg	One tablet once daily on an empty stomach, preferably at bedtime
Complera (RPV/TDF/FTC)	Rilpivirine 25 mg/tenofovir disoproxil fumarate 300 mg/emtricitabine 200 mg	One tablet once daily with a meal
Delstrigo (DOR/TDF/3TC)	Doravirine 100 mg/tenofovir disoproxil fumarate 300 mg/lamivudine 300 mg	One tablet once daily
Odefsey (RPV/TAF/FTC)	Rilpivirine 25 mg/tenofovir alafenamide 25 mg/emtricitabine 200 mg	One tablet once daily with a meal
Symfi (EFV/TDF/3TC)	Efavirenz 600 mg/tenofovir disoproxil fumarate 300 mg/lamivudine 300 mg	One tablet once daily on an empty stomach, preferably at bedtime
Symfi Lo (EFV/TDF/3TC)	Efavirenz 400 mg/tenofovir disoproxil fumarate 300 mg/lamivudine 300 mg	One tablet once daily on an empty stomach, preferably at bedtime
INSTI plus One NNRTI		
Juluca (DTG/RPV)	Dolutegravir 50 mg/rilpivirine 25 mg	One tablet once daily with a meal

^a For dose adjustments in patients with renal or hepatic insufficiency, see [Appendix B, Table 10](#). When no food restriction is listed, the STR can be taken with or without food.

Key: 3TC = lamivudine; ABC = abacavir; ARV = antiretroviral; BIC = bictegravir; DOR = doravirine; DRV = darunavir; DRV/c = darunavir/cobicistat; DTG = dolutegravir; EFV = efavirenz; EVG = elvitegravir; EVG/c = elvitegravir/cobicistat; FDA = Food and Drug Administration; FTC = emtricitabine; INSTI = integrase strand transfer inhibitor; NNRTI = non-nucleoside reverse transcriptase inhibitor; NRTI = nucleoside reverse transcriptase inhibitor; PI = protease inhibitor; RPV = rilpivirine; STR = single-tablet regimen; TAF = tenofovir alafenamide; TDF = tenofovir disoproxil fumarate