

Special Populations: Hepatitis C Virus/HIV Coinfection

(Last updated December 30, 2021; last reviewed December 30, 2021)

Panel's Recommendations
<ul style="list-style-type: none">• All pregnant people with HIV should be screened during the current pregnancy for hepatitis C virus (HCV) infection (AIII).<ul style="list-style-type: none">○ HCV screening should be repeated later in pregnancy in persons who initially screen negative for HCV but who have persistent or new risk factors for HCV (e.g., new or ongoing injection or intranasal substance use) (AIII).• All pregnant people with HIV also should be tested for hepatitis B virus (HBV) infection (see Hepatitis B Virus/HIV Coinfection) (AIII).• Pregnant people with HCV infection who have not already received the hepatitis A virus (HAV) vaccine series should be screened for immunity to HAV (AIII). If they screen negative for HAV antibodies (either immunoglobulin G [IgG] or total antibody [IgG and immunoglobulin M]), they should receive the HAV vaccine series (AIII).• Currently, treatment of HCV during pregnancy is not recommended (unless part of an approved experimental protocol) because of the lack of safety data on the use of HCV direct-acting antiviral agents in persons who are pregnant. If considering initiating HCV treatment in a pregnant person with HCV/HIV coinfection, consultation with an expert in HIV and HCV is strongly recommended (AIII).• Recommendations for antiretroviral therapy (ART) during pregnancy are the same for all people with HIV, including those who have HCV coinfection (AIII).• Pregnant people with HCV/HIV coinfection who are receiving ART should be counseled about the signs and symptoms of liver toxicity, and hepatic transaminases should be assessed 1 month following initiation of ART and at least every 3 months thereafter during pregnancy (BIII).• People with HCV should be strongly considered for HCV treatment with direct-acting antiviral agents postpartum (A).• In people with HCV infection, HCV RNA should be evaluated after delivery to assess for spontaneous clearance of HCV infection, particularly as they are being considered for initiation of HCV therapy postpartum (BII).• HCV/HIV coinfection is not an independent indication for cesarean delivery (see Intrapartum Care for People with HIV) (AIII).• Infants born to people with HCV/HIV coinfection should be evaluated for HCV infection (AIII). Decisions regarding the specific type of assays to use for HCV screening in children and the timing of those assays should be made after consultation with an expert in pediatric HCV infection (AIII).
<p>Rating of Recommendations: A = Strong; B = Moderate; C = Optional</p> <p>Rating of Evidence: I = One or more randomized trials with clinical outcomes and/or validated laboratory endpoints; II = One or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion</p>

The management of hepatitis C virus/HIV coinfection in pregnancy is complex, and none of the approved HCV direct-acting antivirals (DAAs) have been evaluated fully for use in persons who are

pregnant; thus, consultation with an expert in HIV and HCV infection **is strongly recommended** when managing HCV during pregnancy.

For additional information on HCV and HIV, see [Hepatitis C Virus](#) in the [Pediatric Opportunistic Infection Guidelines](#), [Hepatitis C Virus/HIV Coinfection](#) in the [Adult and Adolescent Antiretroviral Guidelines](#), and [Hepatitis C Virus](#) in the [Adult and Adolescent Opportunistic Infection Guidelines](#). The American Association for the Study of Liver Diseases (AASLD), the Infectious Diseases Society of America (IDSA), and the International Antiviral Society–USA maintain updated information about treating patients with HCV/HIV coinfection. The guidelines are available online at HCVguidelines.org.

Screening

All pregnant people with HIV should be screened at entry into general HIV care and during each pregnancy for HCV infection, unless they are known to have HCV/HIV coinfection. **If they are known to have a history of HCV infection, consultation with an expert is recommended for follow-up testing and/or referral for treatment postpartum, as appropriate.**

The primary reasons for HCV testing during pregnancy are—

- To identify pregnant persons with HCV/HIV coinfection at a time when they are engaged with the health care system, so that HCV treatment can be offered after delivery (ideally before a subsequent pregnancy);
- To monitor for HCV-related hepatotoxicity, which has been associated with the use of antiretroviral (ARV) drugs in women with HCV/HIV coinfection;¹
- To monitor for preterm birth, which has been associated with HCV/HIV coinfection in pregnant women;²⁻⁵ and
- To ensure appropriate follow-up and evaluation of infants who were exposed to HCV.

The observed prevalence for HCV infection was 2% to 12% in European cohorts of pregnant women with HIV⁴ and 3.8% among women with HIV in New York State.⁶ Although data about secular trends in HCV among women with HIV in the United States are limited, the prevalence of HCV among women of childbearing age and children aged <2 years in the general population has increased substantially in recent years, partly because of the ongoing opioid epidemic.^{5,7-13}

The Society for Maternal–Fetal Medicine and the American College of Obstetricians and Gynecologists recommend repeating HCV testing later in pregnancy for individuals who initially screen negative for HCV but who have persistent risk factors for HCV or who develop new risk factors for HCV infection (e.g., new or ongoing use of injected or intranasal substance use).¹⁴ The partners of all people with HCV/HIV coinfection should be referred for both HIV and hepatitis counseling and testing to prevent the sexual transmission of HIV and HCV; however, HCV is transmitted infrequently via heterosexual sex. People who do not share injection equipment have a very low risk of horizontal transmission of HCV. Partners who do not have HIV infection should be counseled about the benefits of starting oral pre-exposure prophylaxis (PrEP) to prevent HIV acquisition (see [Pregnancy Counseling and Care for Persons of Childbearing Age with HIV](#)).

Newly available DAAs have improved HCV therapy dramatically; it is now possible to cure HCV infection in most patients.¹⁵ Current HCV treatment guidelines recommend therapy for nearly all patients with HCV infection.¹⁵ However, the management of HCV/HIV coinfection during pregnancy is

complex. A Phase 1 study is now evaluating the safety and pharmacokinetics (PKs) of ledipasvir/sofosbuvir in pregnancy.¹⁶ Safety data of DAAs in pregnancy are still limited. Ribavirin, although it is no longer commonly used for the treatment of HCV, is contraindicated in pregnancy.¹⁷ If considering HCV treatment for a pregnant person, consultation with an expert in HIV and HCV is strongly recommended.

Screening for chronic HCV infection using a sensitive immunoassay for HCV antibodies is recommended for all individuals with HIV, including those who are pregnant. All pregnant people in the United States should be screened for HCV at each pregnancy, except in settings where the prevalence of HCV infection is <0.1%.^{15,18,19} False-negative anti-HCV immunoassay results can occur in individuals with HIV, but this is uncommon with the more sensitive immunoassays. If HCV infection is suspected despite a negative HCV antibody screen, a commercially available diagnostic quantitative plasma HCV RNA assay can be performed.^{20,21} Individuals who have a positive HCV antibody test should undergo confirmatory testing for HCV RNA with this quantitative assay. Many laboratories now perform reflex RNA testing for individuals who test positive for HCV antibodies. Pregnant people also should be tested for HCV RNA when they have indeterminate or negative serologic test results for HCV but are suspected of having HCV infection because of elevated aminotransaminase levels or risk behaviors (e.g., a history of injection drug use).²²

Because of the added risk of hepatic decompensation from acute infection with any viral hepatitis, people with HCV infection also should be screened for both hepatitis A virus (HAV) and hepatitis B virus (HBV). People with chronic HCV infection who have not already received the HAV vaccine series should be screened for immunity to HAV (either immunoglobulin G [IgG] alone or IgG and immunoglobulin M together). If they screen negative for HAV antibodies, they should receive the HAV vaccine series. In patients with CD4 T lymphocyte (CD4+) cell counts <200 cells/mm³, antibody responses to the HAV vaccine should be assessed 1 month after the patient completes the vaccination series; those who are HAV antibody IgG negative should be revaccinated when the CD4 count is >200 cells/mm³.^{23,24} People with HCV/HIV coinfection who screen negative for HBV and lack HBV immunity (i.e., they are hepatitis B surface antigen [HBsAg] negative, hepatitis B core antibody negative, and hepatitis B surface antibody negative [HBsAb]) should receive the HBV vaccine series. Women with HCV/HIV coinfection who are HBsAb negative despite receiving the HBV vaccine series may benefit from revaccination (see [Hepatitis B Virus/HIV Coinfection](#)).²⁵

Impact of HCV/HIV Coinfection on Progression and Perinatal Transmission of Both Viruses

Although the HCV viral load tends to peak in the third trimester, pregnancy does not appear to influence the course of HCV infection clinically. People with chronic HCV generally do well during pregnancy, provided that they have not progressed to decompensated cirrhosis.^{26,27}

Hepatitis C Virus Transmission to the Infant

About 6% of infants born to women with HCV acquire HCV infection.²¹ In most studies of women with HCV/HIV coinfection who are not receiving treatment for either infection, the incidence of perinatal HCV transmission is approximately twofold higher among women with HCV/HIV coinfection (7% to 20% transmission risk) than among women with HCV mono-infection.²⁸⁻³² These higher transmission rates likely are related to the higher levels of HCV viremia observed in patients with HCV/HIV coinfection and/or on other HIV-related impacts on HCV disease activity.^{3,33} Early and sustained control of HIV viremia with ART, however, could reduce the risk of HCV transmission to infants.^{27,34-36} A

European study of perinatal HCV transmission found that the use of effective ART for HIV was associated with a strong trend toward reduced rates of HCV transmission (odds ratio [OR] 0.26; 95% confidence interval [CI], 0.07–1.01).³⁴ In an Italian cohort, HCV transmission occurred in 9% of infants born to women with HCV/HIV coinfection, most of whom were on ART. No HCV transmissions occurred in infants born to women with HCV viral loads of <5 log IU/mL.³

HIV Transmission to the Infant

In the absence of ART, maternal HCV/HIV coinfection can increase the risk of perinatal HIV transmission.^{37,38} The risk of perinatal HIV transmission can be reduced in pregnant people with HCV/HIV coinfection by following the standard recommendations for ART for all women with HIV.

Impact of Hepatitis C Virus on HIV Management

Data are limited on the optimal management of pregnant people with HCV/HIV coinfection. Recommendations on the use of ART during pregnancy for treating HIV and preventing perinatal HIV transmission are the same for people with HCV/HIV coinfection as for those with HIV mono-infection (see [General Principles Regarding Use of Antiretroviral Drugs During Pregnancy](#)). In one Canadian study, HCV/HIV coinfection was associated with an increased risk of HIV viral load **increases near delivery** among women who were on previously effective ART. Although the authors suggest that additional factors (e.g., adherence) may have played a role, these findings support the need to follow recommendations for HIV RNA monitoring during pregnancy.³⁹

Hepatitis C Virus–Specific Therapy in Pregnancy

Several DAA regimens have been approved for the treatment of HCV. At present, all currently available DAAs lack sufficient safety data to be recommended for use during pregnancy, but general considerations for treatment are presented in this section.

When determining the optimal regimen for an individual patient, clinicians must consider many factors, including HCV genotype, prior treatment experience, and stage of liver disease (e.g., compensated or decompensated cirrhosis). The following main classes of DAAs are currently available in the United States:

- NS5A inhibitors: elbasvir, ledipasvir, pibrentasvir, velpatasvir
- NS5B nucleoside polymerase inhibitors: sofosbuvir
- NS3/4A protease inhibitors (PIs): glecaprevir, grazoprevir, voxilaprevir

In the past, most anti-HCV therapy included both interferon and ribavirin. Interferons are not recommended for use in pregnancy because they are abortifacient at high doses in monkeys and have direct antigrowth and antiproliferative effects.⁴⁰ **Pegylated interferon is used rarely for treatment of HCV.** DAA regimens with ribavirin **are indicated for certain regimens and patient populations.** Any treatment regimens that include ribavirin are **contraindicated** for use during pregnancy because of the teratogenic and embryocidal effects observed in all animal species exposed to ribavirin. Ribavirin-associated defects in animals include limb abnormalities, craniofacial defects, anencephaly, and anophthalmia. The risk of teratogenicity persists for up to 6 months following ribavirin cessation and also applies to pregnancies of partners of men taking ribavirin.¹⁵

DAAs are not yet recommended for use in pregnancy because of the lack of PK and safety data; [one small PK study](#) investigating the use of ledipasvir/sofosbuvir in pregnant women with HCV alone demonstrated 100% virologic suppression and no safety concerns. Similarly, a small case series of 15 pregnant women treated with ledipasvir/sofosbuvir reported 100% virologic suppression at 12 weeks and no early safety concerns in the women or their infants.⁴¹ **An open-label, Phase 1 study of ledipasvir-sofosbuvir started between 23 and 24 weeks' gestation in eight women with genotype 1 HCV infection showed that ledipasvir and sofosbuvir exposures were similar in the pregnant women versus the nonpregnant reference group and the drug combination was safe.**¹⁶ Pregnant people with HCV/HIV coinfection should be started on HCV treatment with DAAs postpartum.⁴² Drug interactions exist between the DAA anti-HCV drugs and ARV drugs that may produce clinically significant changes in serum levels of both ARV drugs and anti-HCV medications. For detailed information on the interactions between ARV drugs and anti-HCV drugs, see the [Adult and Adolescent Antiretroviral Guidelines](#), the [Adult and Adolescent Opportunistic Infection Guidelines](#), [HCVGuidelines.org](#), and the [HEP Drug Interaction Checker](#).

Monitoring People with HCV/HIV Coinfection During Pregnancy

Hepatic enzyme levels can increase after ART is initiated in people with HCV/HIV coinfection—particularly in those with low CD4 counts at treatment initiation—as a result of an immune-mediated flare in HCV disease triggered by immune reconstitution with ART. In patients with HIV, HCV coinfection may increase the hepatotoxic risk of certain ARV agents, specifically PIs and nevirapine. HCV monoinfection may increase the risk of intrahepatic cholestasis of pregnancy;⁴³ this risk also is higher among individuals with HCV/HIV coinfection than among individuals with HIV infection alone.⁴ Pregnant patients with HCV/HIV coinfection should be counseled about the signs and symptoms of liver toxicity, and transaminase levels should be assessed 1 month after initiating ART and then every 3 months. If hepatic toxicity occurs, a clinician may need to consider initiating a less hepatotoxic drug regimen, and, if clinical symptoms or significant elevations of transaminases occur, drugs may need to be discontinued temporarily. Differentiating between drug toxicity and a flare of HCV disease that is associated with immune reconstitution can be difficult; therefore, consulting an expert in HCV/HIV coinfection is recommended.

HCV RNA levels can fluctuate during pregnancy and postpartum, with frequent increases in HCV RNA levels during pregnancy followed by a drop in the postpartum period.⁴⁴ Spontaneous clearance of HCV can occur postpartum.⁴⁴⁻⁴⁷ As a result, the AASLD and the IDSA recommend that women have their HCV RNA re-evaluated after delivery, particularly if they are being assessed for initiation of therapy with DAA.⁴²

Rates of preterm delivery are high among individuals with HCV/HIV coinfection. In an Italian cohort of mostly ART-treated women with HCV/HIV coinfection, preterm delivery occurred in 41% of women overall. The rate of preterm delivery was not significantly different among women with lower or higher HCV RNA levels: 29% among women with HCV RNA <5 log IU/mL and 43% among women with HCV RNA >5 log IU/mL. However, women with preterm delivery had significantly higher levels of HCV RNA than those who delivered at term.³ A study of 4,236 pregnant women with HIV reported a higher risk of preterm delivery in women with HCV coinfection (OR 3.0; 95% CI, 1.6–5.7) than in women with HIV alone.⁴ **A study of 339 HIV/HCV coinfecting pregnant women from Spain demonstrated a 50% rate of preterm delivery.**³²

Infants born to women with HCV also were more likely to have low birth weights (defined as weighing <2,500 g) than those born to women without HCV (23 vs. 8%, $P < 0.01$).⁵

HCV infection in pregnancy may be associated with increased risks for gestational diabetes, small-for-gestational-age infants, and low birth weight infants.^{9,48} Although no obstetric guidelines currently suggest that persons with HCV infection should be monitored more frequently for diabetes, preterm birth, or fetal growth during pregnancy,⁴⁹ knowledge of these increased risks may inform clinical care.¹⁴

Mode of Delivery

The majority of studies of scheduled cesarean delivery in women with HCV infection (with or without HIV coinfection) have found that the procedure does not reduce the risk of perinatal HCV transmission.^{34,50-52} Thus, the general recommendations for mode of delivery are the same for people with HCV/HIV coinfection as for those with HIV infection alone (see [Intrapartum Care for People with HIV](#)).

Evaluation of Infants Exposed to Hepatitis C Virus

Infants born to people with HCV/HIV coinfection should be assessed for chronic HCV infection. An HCV antibody test should be performed after age 18 months, when the maternal anti-HCV antibody level has waned.⁵³ Sensitivity of HCV RNA testing is low at birth, and viremia can be intermittent or infection may resolve spontaneously,^{9,54,55} thus, HCV RNA testing should not be performed before age 2 months, and a single negative test is not conclusive evidence of lack of infection.⁵⁶ Uptake of HCV testing is very low for infants who were exposed to HCV;⁵⁷ therefore, it is important for providers to counsel patients about the need for pediatric follow-up and testing during the first few years of life.^{5,58-61} The [Pediatric Opportunistic Infection Guidelines](#) provide further details about the diagnostic evaluation of infants who were exposed to HCV.

References

1. Sibiude J, Warszawski J, Tubiana R, et al. Liver enzyme elevation in pregnant women receiving antiretroviral therapy in the ANRS-French Perinatal Cohort. *J Acquir Immune Defic Syndr*. 2019;81(1):83-94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30702449>.
2. Huang QT, Huang Q, Zhong M, et al. Chronic hepatitis C virus infection is associated with increased risk of preterm birth: a meta-analysis of observational studies. *J Viral Hepat*. 2015;22(12):1033-1042. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26081198>.
3. Baroncelli S, Pirillo MF, Amici R, et al. HCV-HIV coinfecting pregnant women: data from a multicentre study in Italy. *Infection*. 2016;44(2):235-242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26507133>.
4. Benhammou V, Tubiana R, Matheron S, et al. HBV or HCV coinfection in HIV-1-infected pregnant women in France: prevalence and pregnancy outcomes. *J Acquir Immune Defic Syndr*. 2018;77(5):439-450. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29287028>.
5. Chappell CA, Hillier SL, Crowe D, Meyn LA, Bogen DL, Krans EE. Hepatitis C virus screening among children exposed during pregnancy. *Pediatrics*. 2018;141(6):e20173273. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29720535>.
6. Ghazaryan L, Smith L, Parker M, et al. Hepatitis C seroprevalence among HIV-infected childbearing women in New York state in 2006. *Matern Child Health J*. 2016;20(3):550-555. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26520159>.
7. Koneru A, Nelson N, Hariri S, et al. Increased hepatitis C virus (HCV) detection in women of childbearing age and potential risk for vertical transmission—United States and Kentucky, 2011–2014. *MMWR Morb Mortal Wkly Rep*. 2016;65(28):705-710. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27442053>.
8. Ly KN, Jiles RB, Teshale EH, Foster MA, Pesano RL, Holmberg SD. Hepatitis C virus infection among reproductive-aged women and children in the United States, 2006 to 2014. *Ann Intern Med*. 2017;166(11):775-782. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28492929>.
9. Barritt AS, 4th, Jhaveri R. Treatment of hepatitis C during pregnancy—weighing the risks and benefits in contrast to HIV. *Curr HIV/AIDS Rep*. 2018;15(2):155-161. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29470782>.
10. Salemi JL, Spooner KK, Mejia de Grubb MC, Aggarwal A, Matas JL, Salihu HM. National trends of hepatitis B and C during pregnancy across sociodemographic, behavioral, and clinical factors, United States, 1998–2011. *J Med Virol*. 2017;89(6):1025-1032. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27805270>.
11. Patrick SW, Bauer AM, Warren MD, Jones TF, Wester C. Hepatitis C virus infection among women giving birth—Tennessee and United States, 2009–2014. *MMWR Morb Mortal Wkly Rep*. 2017;66(18):470-473. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28493860>.

12. Schillie SF, Canary L, Koneru A, et al. Hepatitis C virus in women of childbearing age, pregnant women, and children. *Am J Prev Med.* 2018;55(5):633-641. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30342628>.
13. Ryerson AB, Schillie S, Barker LK, Kupronis BA, Wester C. Vital Signs: Newly Reported acute and chronic hepatitis C cases—United States, 2009–2018. *MMWR Morb Mortal Wkly Rep.* 2020;69(14):399-404. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32271725>.
14. Society for Maternal-Fetal Medicine, Hughes BL, Page CM, Kuller JA. Hepatitis C in pregnancy: screening, treatment, and management. *Am J Obstet Gynecol.* 2017;217(5):B2-B12. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28782502>.
15. American Association for the Study of Liver Diseases, Infectious Diseases Society of America. HCV guidance: recommendations for testing, managing, and treating hepatitis C. 2020. Available at: <http://www.hcvguidelines.org>.
16. Chappell CA, Scarsi KK, Kirby BJ, et al. Ledipasvir plus sofosbuvir in pregnant women with hepatitis C virus infection: a phase 1 pharmacokinetic study. *Lancet Microbe.* 2020;1(5):e200-e208. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32939459>.
17. Spera AM, Eldin TK, Tosone G, Orlando R. Antiviral therapy for hepatitis C: Has anything changed for pregnant/lactating women? *World J Hepatol.* 2016;8(12):557-565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27134703>.
18. Schillie S, Wester C, Osborne M, Wesolowski L, Ryerson AB. CDC recommendations for hepatitis C screening among adults—United States. *MMWR Recomm Rep* 2020;69(No. RR-2):1–17. Available at: <https://www.cdc.gov/mmwr/volumes/69/rr/rr6902a1.htm>.
19. U.S. Preventive Services Task Force. Final recommendation statement: hepatitis C virus infection in adolescents and adults: screening. 2020. Available at: <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-c-screening>
20. Alter MJ, Kuhnert WL, Finelli L, Centers for Disease Control and Prevention. Guidelines for laboratory testing and result reporting of antibody to hepatitis C virus. Centers for Disease Control and Prevention. *MMWR Recomm Rep.* 2003;52(RR-3):1-13, 15; quiz CE11-14. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12585742>.
21. Centers for Disease Control and Prevention. Hepatitis C Questions and answers for health professionals. 2019. Available at: <https://www.cdc.gov/hepatitis/hcv/hcvfaq.htm>.
22. Centers for Disease Control and Prevention. Viral hepatitis—hepatitis C information. 2019. Available at: <http://www.cdc.gov/hepatitis/hcv>.
23. Panel on Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents with HIV. Table 1. Prophylaxis to prevent first episode of opportunistic disease. 2019. Available at: <https://clinicalinfo.hiv.gov/en/guidelines/adult-and-adolescent-opportunistic-infection/table-1-prophylaxis-prevent-first-episode>.
24. Panel on Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents with HIV. Hepatitis B virus. 2019. Available at:

<https://clinicalinfo.hiv.gov/en/guidelines/adult-and-adolescent-opportunistic-infection/hepatitis-b-virus-infection?view=full>

25. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents—Hepatitis C virus infection. 2021. Available at: <https://clinicalinfo.hiv.gov/en/guidelines/adult-and-adolescent-opportunistic-infection/hepatitis-c-virus-infection>.
26. Sookoian S. Effect of pregnancy on pre-existing liver disease: chronic viral hepatitis. *Ann Hepatol*. 2006;5(3):190-197. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17060881>.
27. Benova L, Mohamoud YA, Calvert C, Abu-Raddad LJ. Vertical transmission of hepatitis C virus: systematic review and meta-analysis. *Clin Infect Dis*. 2014;59(6):765-773. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24928290>.
28. Tovo PA, Palomba E, Ferraris G, et al. Increased risk of maternal–infant hepatitis C virus transmission for women coinfecting with human immunodeficiency virus type 1. Italian Study Group for HCV Infection in Children. *Clin Infect Dis*. 1997;25(5):1121-1124. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9402369>.
29. Gibb DM, Goodall RL, Dunn DT, et al. Mother-to-child transmission of hepatitis C virus: evidence for preventable peripartum transmission. *Lancet*. 2000;356(9233):904-907. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11036896>.
30. Mast EE, Hwang LY, Seto DS, et al. Risk factors for perinatal transmission of hepatitis C virus (HCV) and the natural history of HCV infection acquired in infancy. *J Infect Dis*. 2005;192(11):1880-1889. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16267758>.
31. Alter MJ. Epidemiology of viral hepatitis and HIV co-infection. *J Hepatol*. 2006;44(1 Suppl):S6-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16352363>.
32. Dominguez-Rodriguez S, Prieto L, Fernandez McPhee C, et al. Perinatal HCV Transmission rate in HIV/HCV coinfecting women with access to ART in Madrid, Spain. *PLoS One*. 2020;15(4):e0230109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32271775>.
33. Polis CB, Shah SN, Johnson KE, Gupta A. Impact of maternal HIV coinfection on the vertical transmission of hepatitis C virus: a meta-analysis. *Clin Infect Dis*. 2007;44(8):1123-1131. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17366462>.
34. European Paediatric Hepatitis C Virus Network. A significant sex—but not elective cesarean section—effect on mother-to-child transmission of hepatitis C virus infection. *J Infect Dis*. 2005;192(11):1872-1879. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16267757>.
35. Checa Cabo CA, Stoszek SJ, Quarleri J, et al. Mother-to-child transmission of hepatitis C virus (HCV) among HIV/HCV-coinfecting women. *J Ped Infect Dis*. 2013;2(2):126-135. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26199724>.
36. Conte D, Fraquelli M, Prati D, Colucci A, Minola E. Prevalence and clinical course of chronic hepatitis C virus (HCV) infection and rate of HCV vertical transmission in a cohort of 15,250

- pregnant women. *Hepatology*. 2000;31(3):751-755. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10706568>.
37. Hershov RC, Riester KA, Lew J, et al. Increased vertical transmission of human immunodeficiency virus from hepatitis C virus-coinfected mothers. Women and Infants Transmission Study. *J Infect Dis*. 1997;176(2):414-420. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9237706>.
 38. England K, Thorne C, Newell ML. Vertically acquired paediatric coinfection with HIV and hepatitis C virus. *Lancet Infect Dis*. 2006;6(2):83-90. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16439328>.
 39. Boucoiran I, Albert AYK, Tulloch K, et al. Human immunodeficiency virus viral load rebound near delivery in previously suppressed, combination antiretroviral therapy–treated pregnant women. *Obstet Gynecol*. 2017;130(3):497-501. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28796673>.
 40. Boskovic R, Wide R, Wolpin J, Bauer DJ, Koren G. The reproductive effects of beta interferon therapy in pregnancy: a longitudinal cohort. *Neurology*. 2005;65(6):807-811. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16186517>.
 41. Yattoo GN. Treatment of chronic hepatitis C with ledipasvir/sofosbuvir combination during pregnancy [Abstract]. *Hepatol Int*. 2018;12(2):S292-S293. Available at: <https://www.hcvguidelines.org/references/yattoo-2018>.
 42. American Association for the Study of Liver Diseases, Infectious Diseases Society of America. HCV in pregnancy 2020. Available at: <https://www.hcvguidelines.org/unique-populations/pregnancy>.
 43. Wijarnpreecha K, Thongprayoon C, Sanguankeo A, Upala S, Ungprasert P, Cheungpasitporn W. Hepatitis C infection and intrahepatic cholestasis of pregnancy: a systematic review and meta-analysis. *Clin Res Hepatol Gastroenterol*. 2017;41(1):39-45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27542514>.
 44. Lin HH, Kao JH. Hepatitis C virus load during pregnancy and puerperium. *BJOG*. 2000;107(12):1503-1506. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11192107>.
 45. Hattori Y, Orito E, Ohno T, et al. Loss of hepatitis C virus RNA after parturition in female patients with chronic HCV infection. *J Med Virol*. 2003;71(2):205-211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12938194>.
 46. Honegger JR, Kim S, Price AA, et al. Loss of immune escape mutations during persistent HCV infection in pregnancy enhances replication of vertically transmitted viruses. *Nat Med*. 2013;19(11):1529-1533. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24162814>.
 47. Hashem M, Jhaveri R, Saleh DA, et al. Spontaneous viral load decline and subsequent clearance of chronic hepatitis C Virus in postpartum women correlates with favorable interleukin-28B gene allele. *Clin Infect Dis*. 2017;65(6):999-1005. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28903504>.

48. Pergam SA, Wang CC, Gardella CM, Sandison TG, Phipps WT, Hawes SE. Pregnancy complications associated with hepatitis C: data from a 2003–2005 Washington state birth cohort. *Am J Obstet Gynecol*. 2008;199(1):38 e31-39. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18486089>.
49. American College of Obstetricians Gynecologists. ACOG practice bulletin No. 86: viral hepatitis in pregnancy. *Obstet Gynecol*. 2007;110(4):941-956. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17906043>.
50. Ghamar Chehreh ME, Tabatabaei SV, Khazanehdari S, Alavian SM. Effect of cesarean section on the risk of perinatal transmission of hepatitis C virus from HCV-RNA+/HIV- mothers: a meta-analysis. *Arch Gynecol Obstet*. 2011;283(2):255-260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20652289>.
51. Marine-Barjoan E, Berrebi A, Giordanengo V, et al. HCV/HIV co-infection, HCV viral load and mode of delivery: risk factors for mother-to-child transmission of hepatitis C virus? *AIDS*. 2007;21(13):1811-1815. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17690581>.
52. McMenamin MB, Jackson AD, Lambert J, et al. Obstetric management of hepatitis C-positive mothers: analysis of vertical transmission in 559 mother–infant pairs. *Am J Obstet Gynecol*. 2008;199(3):315 e311-315. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18771997>.
53. Bal A, Petrova A. Single clinical practice's report of testing initiation, antibody clearance, and transmission of hepatitis C virus (HCV) in infants of chronically HCV-infected mothers. *Open Forum Infect Dis*. 2016;3(1):ofw021. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26985444>.
54. Mack CL, Gonzalez-Peralta RP, Gupta N, et al. NASPGHAN practice guidelines: Diagnosis and management of hepatitis C infection in infants, children, and adolescents. *J Pediatr Gastroenterol Nutr*. 2012;54(6):838-855. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22487950>.
55. Bernstein HB, Dunkelberg JC, Leslie KK. Hepatitis C in pregnancy in the era of direct-acting antiviral treatment: potential benefits of universal screening and antepartum therapy. *Clin Obstet Gynecol*. 2018;61(1):146-156. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29351151>.
56. Polywka S, Pembrey L, Tovo PA, Newell ML. Accuracy of HCV-RNA PCR tests for diagnosis or exclusion of vertically acquired HCV infection. *J Med Virol*. 2006;78(2):305-310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16372293>.
57. Lopata SM, McNeer E, Dudley JA, et al. Hepatitis C testing among perinatally exposed infants. *Pediatrics*. 2020;145(3):e20192482. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32060140>.
58. Kuncio DE, Newbern EC, Johnson CC, Viner KM. Failure to test and identify perinatally infected children born to hepatitis C virus-infected women. *Clin Infect Dis*. 2016;62(8):980-985. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26797211>.
59. Watts T, Stockman L, Martin J, Guilfoyle S, Vergeront JM. Increased risk for mother-to-infant transmission of hepatitis C virus among medicaid recipients—Wisconsin, 2011–2015. *MMWR*

Morb Mortal Wkly Rep. 2017;66(42):1136-1139. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29072864>.

60. Towers CV, Fortner KB. Infant follow-up postdelivery from a hepatitis C viral load positive mother. *J Matern Fetal Neonatal Med.* 2019;32(19):3303-3305. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29587561>.
61. Gowda C, Smith S, Crim L, Moyer K, Sanchez PJ, Honegger JR. Nucleic acid testing for diagnosis of perinatally-acquired hepatitis C Virus Infection in early infancy. *Clin Infect Dis.* 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32640018>.