

## Teratogenicity

Updated: January 31, 2023

Reviewed: January 31, 2023

Panel's Recommendations
<ul style="list-style-type: none"><li>All cases of antiretroviral (ARV) drug exposure during pregnancy should be reported to the <a href="#">Antiretroviral Pregnancy Registry (AIII)</a>.</li><li>Based on multiple studies indicating no difference in rates of total birth defects for first-trimester exposure compared with later ARV drug exposures, people can be counseled that ARV drugs during pregnancy generally do not increase the risk of birth defects (BIII). Providers should be aware that data on the risks of birth defects for many ARV drugs are limited and evolving (see <a href="#">Appendix B: Supplement: Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy</a>).</li><li>All pregnant people with HIV should initiate antiretroviral therapy (ART) as early in pregnancy as possible (AI). Pregnant people with HIV should not delay initiating ART due to concerns about teratogenicity with first trimester exposure (AIII).<ul style="list-style-type: none"><li>The Panel on Treatment of HIV During Pregnancy and Prevention of Perinatal Transmission emphasizes the importance of counseling and informed decision-making regarding all ARV regimens for people with HIV (AIII). For additional information, see <a href="#">Appendix C: Antiretroviral Counseling Guide for Health Care Providers</a>.</li></ul></li><li>Clinicians should discuss <a href="#">future reproductive plans and timing, as well as the risks and benefits of conceiving on specific ARV medications</a>, and the use of appropriate contraceptive options to prevent unintended pregnancies (AIII). See <a href="#">Introduction to the Selection of Antiretroviral Drugs In Pregnancy</a>, People with HIV Who are Trying to Conceive, and <a href="#">Table 7</a>.</li></ul>
<p><i>Rating of Recommendations: A = Strong; B = Moderate; C = Optional</i></p> <p><i>Rating of Evidence: I = One or more randomized trials with clinical outcomes and/or validated laboratory endpoints; II = One or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion</i></p>

### ***Antiretroviral Pregnancy Registry Reporting***

Health care providers who are caring for pregnant people with HIV are advised strongly to report instances of prenatal exposure to antiretroviral (ARV) drugs (either single-drug exposure or exposure to a combination of ARV drugs) to the [Antiretroviral Pregnancy Registry](#) as early in pregnancy as possible. The purpose of the Antiretroviral Pregnancy Registry is to detect any major teratogenic effect involving any of the registry drugs to which pregnant people are exposed. Registry data are used to supplement animal toxicology studies and assist clinicians in weighing the potential risks and benefits of treatment for individual patients. The Antiretroviral Pregnancy Registry is a collaborative project of pharmaceutical manufacturers with an advisory committee that includes a teratologist; an infectious disease specialist; an epidemiologist; a biostatistician; and a group of obstetric, maternal–fetal medicine, and pediatric providers. This prospective registry does not use patient names, and registry staff obtain birth outcome follow-up information from the reporting health care provider.

Referrals should be directed to—

Antiretroviral Pregnancy Registry  
Research Park  
301 Government Center Drive  
Wilmington, NC 28403  
Telephone: 1-800-258-4263

Fax: 1-800-800-1052

Email: [SM\\_APR@APRegistry.com](mailto:SM_APR@APRegistry.com)

## ***Antiretroviral Drugs and Birth Defects***

The potential harm to the fetus from maternal ingestion of a specific drug depends not only on the drug itself, but also on the dose ingested; the gestational age of the fetus at exposure; the duration of exposure; interactions with other agents to which the fetus is exposed; and, to an unknown extent, the genetic makeup of the mother and fetus. Information regarding the safety of using certain drugs during pregnancy is derived from multiple sources, including animal reproductive/developmental toxicity data, anecdotal experience, registry data, randomized clinical trials, and observational studies.

Drug choice should be individualized and discussed with the people who are pregnant or are trying to conceive before treatment begins. Clinicians also must consider available data from preclinical and clinical testing of the individual drugs. Preclinical data include results of *in vitro* and animal *in vivo* screening tests for carcinogenicity, clastogenicity/mutagenicity, and reproductive and teratogenic effects. However, the predictive value of such tests for adverse effects in humans is unknown. When considering whether a pregnant person should continue an effective antiretroviral regimen when they present in early pregnancy, the potential risk of viral rebound with switching regimens must be considered, as well as the specific or unknown risks for birth defects of the current drug regimen and stage of gestation.<sup>1</sup> For additional information, see [People With HIV Who Are Taking Antiretroviral Therapy When they Become Pregnant](#).

Data continue to be collected on the placental passage, pharmacokinetics, and safety of U.S. Food and Drug Administration (FDA)–approved ARV drugs administered during pregnancy, in addition to data on the long-term safety in infants who were exposed to these drugs *in utero*. However, the data remain somewhat limited, especially for newer drugs (see [Appendix B: Supplement: Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy](#)). The Antiretroviral Pregnancy Registry has predefined analytical methods and criteria for recognizing a potential signal. When analyzing registry data, data on birth outcomes from 200 infants who were exposed to an ARV drug during the first trimester are viewed as sufficient to detect a doubling of the risk of overall birth defects associated with that drug compared to the general population. A cohort of 1,000 is sufficient to detect a 1.5-fold increase in the risk of overall birth defects. The general U.S. population birth defect prevalence is 2.72% as determined by the Metropolitan Atlanta Congenital Defect Program, a population-based tracking system for birth defects.<sup>2</sup> summarizes Antiretroviral Pregnancy Registry risk assessment for individual ARV drugs and points out that risk assessment is Table 8 not available when pregnancy exposures are not reported. Detailed information about Antiretroviral Pregnancy Registry data for individual drugs is available in [Appendix B: Supplement: Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy](#).

**Table 8. Drug-Specific Risk Assessment by the Antiretroviral Pregnancy Registry**

ARV Drug	Level of Risk Assessment	Risk Assessment Outcome
COBI, DRV, d4T, ddI, DTG, EVG, IDV, RAL, RPV, and TAF	Sufficient numbers of first-trimester exposures have been monitored to detect at least a 2-fold increase in the risk of overall birth defects.	No such increases detected

3TC, ABC, ATV, EFV, FTC, LPV/r, NFV, NVP, RTV, TDF, and ZDV	Sufficient numbers of first-trimester exposures have been monitored to detect at least a 1.5-fold increase in the risk of overall birth defects and a twofold increase in the risk of birth defects in cardiovascular and genitourinary systems.	No such increases detected
BIC, CAB, DOR, ETR, FTR, and T-20	Insufficient numbers of exposures reported to assess level of risk	Not available

Key: 3TC = lamivudine; ABC = abacavir; ARV = antiretroviral; ATV = atazanavir; BIC = bictegravir; CAB = cabotegravir; COBI = cobicistat; d4T = stavudine; ddI = didanosine; DOR = doravirine; DRV = darunavir; DTG = dolutegravir; EFV = efavirenz; ETR = etravirine; EVG = elvitegravir; FTC = emtricitabine; FTR = fostemsavir; IDV = indinavir; LPV/r = lopinavir /ritonavir; NFV = nelfinavir; NVP = nevirapine; RAL = raltegravir; RPV = rilpivirine; RTV = ritonavir; T-20 = enfuvirtide; TAF = tenofovir alafenamide; TDF = tenofovir disoproxil fumarate; ZDV = zidovudine

For individual birth defects, the power of the Antiretroviral Pregnancy Registry to find an increased risk will vary depending upon the frequency of the defect in the population. However, data from a larger number of infants are required to detect an increased risk of specific birth defects with lower frequencies of occurrence, with the required number of infants who were exposed to an ARV drug increasing as the frequency of the defect in an unexposed population decreases. Thus, large numbers of cases are required to detect increased risk of rare but serious defects, underscoring the need for providers to report all ARV exposures prospectively to the Antiretroviral Pregnancy Registry.<sup>3</sup>

It is important to consider potential confounding factors in studies of ARV drugs and birth defects. Several factors that are associated with HIV also may increase the risk of birth defects, such as exposure to folate antagonists (e.g., trimethoprim-sulfamethoxazole),<sup>4</sup> nutritional and folate status,<sup>5</sup> and tobacco and alcohol use.<sup>6</sup> Clinicians also should be aware of indication bias, which can occur when a patient’s reason for taking a particular ARV drug is associated with an increased risk of birth defects, such as older age or more advanced disease. Additionally, clinicians should consider all medications used in early pregnancy. According to a 2018 study involving 9,546 pregnant people, 97.1% reported taking at least one medication during their pregnancy.<sup>7</sup> In the last decade, 89.3% of the 290 therapeutics submitted to the FDA between 2010 and 2019 lacked human data related to pregnancy.<sup>8</sup> Thus, it is important to know of any and all medication exposures in pregnancy when evaluating risk of birth defects in relation to ARV drugs.

Several studies of birth defects in fetuses and infants of women who received various ARV regimens during observational studies found no difference in rates of total birth defects between first-trimester drug exposures and later exposures.<sup>9-13</sup> The Antiretroviral Pregnancy Registry conducts a primary analysis of prospective cases of ARV drug exposure during pregnancy provided by health care providers. In the current analysis through January 31, 2022, the prevalence of birth defects was 2.8 per 100 live births among women with a first-trimester exposure to any ARV drug (310 of 10,950 exposures; 95% confidence interval [CI], 2.5–3.2). The prevalence of defects is not significantly different from that seen in women with an initial exposure during the second and/or third trimester (2.9 per 100 live births; prevalence ratio 0.99; 95% CI, 0.84–1.16).<sup>2</sup> Although these studies are reassuring, an increased risk of specific abnormalities—particularly rare abnormalities—would not necessarily be detectable when looking only at the total number of birth defects. Furthermore, risk may be underestimated when defects are ascertained only after live births because this does not include more severe defects that result in stillbirths and terminations. Another limitation is that an increased risk that is associated with a specific ARV drug may be obscured when the analysis unit combines all ARV drugs together. As new drugs are studied in pregnancy, data from animal studies and/or limited and early data from human studies may suggest concerns about specific birth defects,

such as was observed with initial data on a potential association of neural tube defects with periconception use of dolutegravir that, after collection of many additional exposures, was shown to not be increased compared to periconception use of non-dolutegravir regimens.<sup>14</sup> This demonstrates the importance of having data on enough exposures to enable conclusions to be drawn and of reporting such exposures to the Antiretroviral Pregnancy Registry. Additional data and further studies are needed to assess and understand the risk associated with a particular ARV drug. Drug-specific teratogenicity data are summarized in [Appendix B: Supplement: Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy](#).

## References

1. Frange P, Tubiana R, Sibiude J, et al. Rilpivirine in HIV-1-positive women initiating pregnancy: to switch or not to switch? *J Antimicrob Chemother.* 2020;75(5):1324-1331. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32157283>.
2. Antiretroviral Pregnancy Registry Steering Committee. Antiretroviral pregnancy registry international interim report for 1 January 1989–31 January 2022. Wilmington, NC: Registry Coordinating Center; 2022. Available at: <http://www.apregistry.com>.
3. Watts DH. Teratogenicity risk of antiretroviral therapy in pregnancy. *Curr HIV/AIDS Rep.* 2007;4(3):135-140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17883999>.
4. Ford N, Shubber Z, Jao J, Abrams EJ, Frigati L, Mofenson L. Safety of cotrimoxazole in pregnancy: a systematic review and meta-analysis. *J Acquir Immune Defic Syndr.* 2014;66(5):512-521. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24853309>.
5. Jungmann EM, Mercey D, DeRuiter A, et al. Is first trimester exposure to the combination of antiretroviral therapy and folate antagonists a risk factor for congenital abnormalities? *Sex Transm Infect.* 2001;77(6):441-443. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11714944>.
6. Lipshultz SE, Williams PL, Zeldow B, et al. Cardiac effects of in-utero exposure to antiretroviral therapy in HIV-uninfected children born to HIV-infected mothers. *AIDS.* 2015;29(1):91-100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25562493>.
7. Haas DM, Marsh DJ, Dang DT, et al. Prescription and other medication use in pregnancy. *Obstet Gynecol.* 2018;131(5):789-798. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29630018>.
8. Byrne JJ, Saucedo AM, Spong CY. Evaluation of drug labels following the 2015 pregnancy and lactation labeling rule. *JAMA Netw Open.* 2020;3(8):e2015094. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32865574>.
9. Watts DH, Huang S, Culnane M, et al. Birth defects among a cohort of infants born to HIV-infected women on antiretroviral medication. *J Perinat Med.* 2011;39(2):163-170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21142844>.
10. Knapp KM, Brogly SB, Muenz DG, et al. Prevalence of congenital anomalies in infants with in utero exposure to antiretrovirals. *Pediatr Infect Dis J.* 2012;31(2):164-170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21983213>.
11. da Costa TP, Machado ES, et al. Malformations among HIV vertically exposed newborns – results from a Brazilian cohort study. Presented at: 6th IAS Conference on HIV Pathogenesis and Treatment and Prevention; 2011. Rome, Italy.

12. Floridia M, Mastroiacovo P, Tamburrini E, et al. Birth defects in a national cohort of pregnant women with HIV infection in Italy, 2001-2011. *BJOG*. 2013;120(12):1466-1475. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23721372>.
13. Money D, Lee T, O'Brien C, et al. Congenital anomalies following antenatal exposure to dolutegravir: a Canadian surveillance study. *BJOG*. 2019;126(11):1338-1345. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31188522>.
14. Zash R, Holmes L.B., Diseko M., Jacobson D., et al. Update on neural tube defects with antiretroviral exposure in the Tsepamo Study, Botswana,,. Presented at: AIDS 2022; 2022. Montreal, Canada. Available at: <https://programme.aids2022.org/Abstract/Abstract/?abstractid=12759>.