

Cobicistat (COBI, Tybost)

Updated: April 11, 2023

Reviewed: April 11, 2023

Formulations	
<p>Tablet: 150 mg</p> <p>Fixed-Dose Combination (FDC) Tablets</p> <ul style="list-style-type: none"> [Evotaz] Atazanavir 300 mg/cobicistat 150 mg [Genvoya] Elvitegravir 150 mg/cobicistat 150 mg/emtricitabine 200 mg/tenofovir alafenamide 10 mg [Prezcobix] Darunavir 800 mg/cobicistat 150 mg [Stribild] Elvitegravir 150 mg/cobicistat 150 mg/emtricitabine 200 mg/tenofovir disoproxil fumarate 300 mg [Symtuza] Darunavir 800 mg/cobicistat 150 mg/emtricitabine 200 mg/ tenofovir alafenamide 10 mg <p>When using FDC tablets, refer to other sections of Appendix A: Pediatric Antiretroviral Drug Information for information about the individual components of the FDC. See also Appendix A, Table 2. Antiretroviral Fixed-Dose Combination Tablets: Minimum Body Weights and Considerations for Use in Children and Adolescents.</p> <p>For additional information, see Drugs@FDA or DailyMed.</p>	
Dosing Recommendations	Selected Adverse Events
<p>Cobicistat Is a Pharmacokinetic Enhancer</p> <ul style="list-style-type: none"> The only use of cobicistat (COBI) is as a pharmacokinetic (PK) enhancer (boosting agent) for certain protease inhibitors (PIs) and integrase strand transfer inhibitors. COBI is not interchangeable with ritonavir (RTV) and has no antiviral activity. <p>Child and Adolescent (Weighing ≥ 35 kg) and Adult Dose</p> <ul style="list-style-type: none"> COBI 150 mg with atazanavir (ATV) 300 mg administered at the same time with food <p>Child and Adolescent (Weighing ≥ 40 kg) and Adult Dose</p> <ul style="list-style-type: none"> COBI 150 mg with darunavir (DRV) 800 mg administered at the same time with food <p>[Evotaz] Atazanavir/Cobicistat</p> <p><i>Child and Adolescent (Weighing ≥ 35 kg) and Adult Dose</i></p> <ul style="list-style-type: none"> One tablet once daily with food Use in combination with other antiretroviral (ARV) drugs. 	<ul style="list-style-type: none"> COBI is an inhibitor of renal tubular transporters of creatinine. This increases serum creatinine and reduces the estimated glomerular filtration rate, with no change in glomerular function. <p style="text-align: center;">Special Instructions</p> <ul style="list-style-type: none"> COBI 150 mg is not interchangeable with RTV, but it has a PK boosting effect that is comparable to RTV 100 mg. Drug interactions may differ between RTV and COBI, because COBI is a stronger P-glycoprotein inhibitor and lacks some of the induction effects of RTV. Do not administer COBI with RTV or with FDC tablets that contain COBI. COBI is not recommended for use with more than one ARV drug that requires PK enhancement (e.g., elvitegravir used in combination with a PI). Using COBI with PIs other than once-daily ATV 300 mg or DRV 800 mg is not recommended. Patients with a confirmed increase in serum creatinine >0.4 mg/dL from baseline should be closely monitored for renal safety.

<p>[Genvoya] Elvitegravir/Cobicistat/Emtricitabine (FTC)/ Tenofovir Alafenamide (TAF)</p> <p><i>Child (Weighing ≥14 to <25 kg)</i></p> <ul style="list-style-type: none"> Limited data currently exist on the appropriate dose of Genvoya in children ≥14 kg to <25 kg. Studies are currently being conducted to assess the safety and efficacy of a low-dose tablet with elvitegravir (EVG) 90 mg/COBI 90 mg/FTC 120 mg/TAF 6 mg. 	<ul style="list-style-type: none"> When using COBI in combination with TDF, monitor serum creatinine, urine protein, and urine glucose at baseline and every 3 to 6 months while the patient is receiving therapy (see Table 17i. Nephrotoxic Effects). In patients who are at risk of renal impairment, serum phosphate also should be monitored. For information on crushing and cutting tablets, see Information on Crushing and Liquid Drug Formulations (PDF) from Toronto General Hospital.
<p><i>Child and Adolescent (Weighing ≥25 kg) and Adult Dose</i></p> <ul style="list-style-type: none"> One tablet once daily with food 	<p>Metabolism/Elimination</p>
<p>[Prezcobix] Darunavir/Cobicistat</p> <p><i>Child and Adolescent (Weighing ≥40 kg) and Adult Dose</i></p> <ul style="list-style-type: none"> One tablet once daily with food Use in combination with other ARV drugs. 	<ul style="list-style-type: none"> COBI is a strong inhibitor of cytochrome P450 (CYP) 3A4 and a weak inhibitor of CYP2D6.
<p>[Stribild] Elvitegravir/Cobicistat/Emtricitabine/Tenofovir Disoproxil Fumarate (TDF)</p> <p><i>Child and Adolescent (Weighing ≥35 kg) and Adult Dose</i></p> <ul style="list-style-type: none"> One tablet once daily with food The Panel on Antiretroviral Therapy and Medical Management of Children Living With HIV recommends using Stribild only in patients with sexual maturity ratings of 4 or 5. 	<p>Cobicistat Dosing in Patients With Hepatic Impairment</p> <ul style="list-style-type: none"> COBI does not require dose adjustment in patients with mild-to-moderate hepatic impairment. No data are available in patients with severe hepatic impairment. Dosing recommendations for medications that are coadministered with COBI should be followed.¹ Genvoya, Prezcobix, Stribild, and Symtuza are not recommended in patients with severe hepatic impairment.¹ Evotaz is not recommended in patients with any degree of hepatic impairment.
<p>[Symtuza] Darunavir/Cobicistat/Emtricitabine/TAF</p> <p><i>Child and Adolescent (Weighing ≥40 kg) and Adult Dose</i></p> <ul style="list-style-type: none"> One tablet once daily with food 	<p>Cobicistat Dosing in Patients With Renal Impairment</p> <ul style="list-style-type: none"> COBI does not require a dose adjustment in patients with renal impairment, including those with severe renal impairment. Dosing recommendations for medications that are coadministered with COBI should be followed. The use of COBI plus TDF is not recommended in patients with creatinine clearance (CrCl) <70 mL/min. Dose adjustments for TDF are required for patients with CrCl <50 mL/min, and the necessary dose adjustments for TDF when this drug is used with COBI have not been established in this group of patients. Genvoya² is not recommended in patients with estimated CrCl 15 to <30 mL/min, or in patients with estimated CrCl <15 mL/min who are not receiving chronic hemodialysis. Stribild³ should not be initiated in patients with estimated CrCl <70 mL/min and should be discontinued in patients with estimated CrCl <50 mL/min. The dose adjustments required for emtricitabine and TDF in these patients cannot be achieved with an FDC tablet. Symtuza⁴ is not recommended in patients with estimated CrCl <30 mL/min.

Drug Interactions

Additional information about drug interactions is available in the [Adult and Adolescent Antiretroviral Guidelines](#) and the [HIV Drug Interaction Checker](#).

- *Metabolism:* Metabolism of cobicistat (COBI) is mainly via cytochrome P450 (CYP) 3A4 and, to a lesser degree, CYP2D6. COBI is a strong inhibitor of CYP3A4 and a weak inhibitor of CYP2D6. COBI also inhibits breast cancer resistance protein, P-glycoprotein (P-gp), the organic anion transporting polypeptides OATP1B1 and OATP1B3, and multidrug and toxin extrusion 1. Unlike ritonavir, COBI does not demonstrate any enzyme-inducing effects. The potential exists for multiple drug interactions when using COBI. Before COBI is administered, a patient's medication profile should be carefully reviewed for potential interactions and overlapping toxicities with other drugs. Coadministration of medications that induce or inhibit CYP3A4 may respectively decrease or increase exposures of COBI and coformulated antiretroviral medications. Coadministration of medications that are CYP3A4 substrates may result in clinically significant adverse reactions that are severe, life-threatening, or fatal, or may result in loss of therapeutic effect if dependent on conversion to an active metabolite due to CYP3A4 inhibition by COBI.¹
- *Nucleoside reverse transcriptase inhibitors:* COBI is a strong P-gp inhibitor; thus, a dose of tenofovir alafenamide (TAF) 10 mg combined with COBI produces tenofovir (TFV) exposures that are similar to those produced by TAF 25 mg without COBI.⁵ COBI increases plasma TFV exposures by 23% when it is coadministered with TDF; thus, renal safety should be monitored in patients who are receiving this combination.^{1,6}
- *Non-nucleoside reverse transcriptase inhibitors:* Efavirenz, etravirine, and nevirapine **should not be used** with COBI.
- *Protease inhibitors:* Using COBI as a dual booster for elvitegravir (EVG) and darunavir (DRV) has been studied in people with and without HIV, and the evidence is conflicting. When EVG plus COBI plus DRV was administered to people without HIV, the trough concentration (C_{trough}) of EVG was 50% lower than the C_{trough} seen in people who received elvitegravir/cobicistat/emtricitabine/tenofovir disoproxil fumarate (EVG/c/FTC/TDF) without DRV.⁷ When EVG/c/FTC/TAF was administered with DRV to patients with HIV, both DRV and EVG concentrations were comparable to those seen in historic controls.⁸
- *Integrase inhibitors:* In one small study, dolutegravir (DTG) C_{trough} was 107% higher when DTG was administered with darunavir/cobicistat (DRV/c) than when it was administered with darunavir/ritonavir.⁹ Bictegravir (BIC) area under the curve increases 74% when BIC is administered with DRV/c.¹⁰
- *Corticosteroids:* Increased serum concentrations of corticosteroids can occur when corticosteroids and COBI are coadministered; this can lead to clinically significant adrenal suppression. Adrenal suppression occurs regardless of whether the corticosteroids are administered orally or by some other route (e.g., intranasal, inhaled, interlaminar, intraarticular) and regardless of whether the corticosteroids are administered routinely or intermittently. A possible exception is beclomethasone, which appears to be a relatively safe option with inhaled or intranasal administration.^{11,12}

Major Toxicities

- *More common:* Nausea, vomiting, diarrhea, abdominal pain, anorexia
- *Less common (more severe):* New onset renal impairment or worsening of renal impairment when used with **TAF** or TDF; rhabdomyolysis; increased amylase and lipase levels.

Resistance

Not applicable because COBI has no antiviral activity.

Pediatric Use

Approval

COBI is a pharmacokinetic (PK) enhancer of antiretroviral drugs that is available as a single agent or a component of fixed-dose combination products. COBI, as a component of Stribild, is approved by the U.S. Food and Drug Administration (FDA) at the adult dose for use in children and adolescents aged ≥ 12 years and weighing ≥ 35 kg.³ The Panel on Antiretroviral Therapy and Medical Management of Children Living With HIV recommends limiting the use of Stribild to those with a sexual maturity rating of 4 or 5. COBI, as a component of Genvoya, is approved by the FDA at the adult dose for use in children weighing ≥ 25 kg.² The FDA has not approved COBI as a component of Genvoya for use in children < 25 kg, but an ongoing PK, safety, and efficacy study is underway with a low-dose tablet in children weighing ≥ 14 kg to < 25 kg (see the [Elvitegravir](#) section). COBI alone (as Tybost) is approved by the FDA at the adult dose for use in children weighing ≥ 35 kg when used in combination with ATV, and in children weighing ≥ 40 kg when used in combination with DRV.¹ COBI, coformulated with ATV (as Evotaz),¹³ is approved by the FDA at the adult dose for use in children and adolescents weighing ≥ 35 kg. COBI, coformulated with DRV (as Prezcoibix)¹⁴ and as a component of Symtuza,⁴ is approved by the FDA at the adult dose in children and adolescents weighing ≥ 40 kg.

References

1. Cobicistat (Tybost) [package insert]. Food and Drug Administration. 2021. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/203094s016lbl.pdf.
2. Genvoya (elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide) [package insert]. Food and Drug Administration. 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/207561s029lbl.pdf.
3. Stribild (elvitegravir, cobicistat, emtricitabine, tenofovir disoproxil fumarate) [package insert]. Food and Drug Administration. 2021. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/203100s036lbleet.pdf.
4. Symtuza (Darunavir, cobicistat, emtricitabine, and tenofovir alafenamide) [package insert]. Food and Drug Administration. 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/210455s022lbl.pdf.
5. Ramanathan S, Wei X, Custudio J, et al. Pharmacokinetics of a novel EVG/COBI/FTC/GS-7340 single tablet regimen. Abstract O-13. Presented at: 13th International Workshop on Clinical Pharmacology of HIV Therapy; 2012. Barcelona, Spain. Available at: https://www.natap.org/2012/pharm/Pharm_24.htm.
6. Custodio J, Garner W, Jin F, et al. Evaluation of the drug interaction potential between the pharmacokinetic enhancer and tenofovir disoproxil fumarate in healthy subjects. Presented at: 14th International Workshop on Clinical Pharmacology of HIV Therapy; 2013. Amsterdam, The Netherlands.
7. Ramanathan S, Wang H, Szwarcberg J, Kearney BP. Safety/tolerability, pharmacokinetics, and boosting of twice-daily cobicistat administered alone or in combination with darunavir or tipranavir. Abstract abstract P-08. Presented at: 13th International Workshop on Clinical Pharmacology of HIV Therapy; 2012. Barcelona, Spain. Available at: https://www.natap.org/2012/pharm/Pharm_28.htm.
8. Gutierrez-Valencia A, Trujillo-Rodriguez M, Fernandez-Magdaleno T, Espinosa N, Viciano P, Lopez-Cortes LF. Darunavir/cobicistat showing similar effectiveness as darunavir/ritonavir monotherapy despite lower trough concentrations. *J Int AIDS Soc.* 2018;21(2). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29430854>.
9. Gervasoni C, Riva A, Cozzi V, et al. Effects of ritonavir and cobicistat on dolutegravir exposure: when the booster can make the difference. *J Antimicrob Chemother.* 2017;72(6):1842-1844. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28333266>.
10. Zhang H, Custudio J, Wei X, et al. Clinical pharmacology of the HIV integrase strand transfer inhibitor bictegravir. Abstract 40. Presented at: Conference on Retroviruses and Opportunistic Infections; 2017. Seattle, WA. Available at: <https://www.croiconference.org/sessions/clinical-pharmacology-hiv-integrase-strand-transfer-inhibitor-bictegravir>.

11. Saberi P, Phengrasamy T, Nguyen DP. Inhaled corticosteroid use in HIV-positive individuals taking protease inhibitors: a review of pharmacokinetics, case reports and clinical management. *HIV Med.* 2013;14(9):519-529. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23590676>.
12. Boyd SD, Hadigan C, McManus M, et al. Influence of low-dose ritonavir with and without darunavir on the pharmacokinetics and pharmacodynamics of inhaled beclomethasone. *J Acquir Immune Defic Syndr.* 2013;63(3):355-361. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23535292>.
13. Evotaz (atazanavir/cobicistat) [package insert]. Food and Drug Administration. 2020. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/206353s007lbl.pdf.
14. Prezcobix (darunavir/cobicistat) [package insert]. Food and Drug Administration. 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/205395s024lbl.pdf.