

Special Considerations for Antiretroviral Therapy Use in Adolescents with HIV

Updated: Apr. 11, 2022
Reviewed: Apr. 11, 2022

Panel's Recommendations
<ul style="list-style-type: none">• All adolescents with HIV should receive maximally suppressive antiretroviral (ARV) therapy; this is urgent for those who are sexually active, considering pregnancy, or pregnant (AII).• ARV regimen selection should include consideration of the adolescent's individual needs and preferences (AIII). See What to Start and Management of Children Receiving Antiretroviral Therapy for more information.• All adolescents with HIV should be screened for mental health disorders and substance use disorders (AII).• Reproductive health issues—including pregnancy intentions, contraceptive methods, safer sex techniques to prevent transmission of HIV and other sexually transmitted infections, pre-exposure prophylaxis for partners, pregnancy planning, and preconception care—should be discussed regularly (AI).• Providers should be aware of potential interactions between specific ARV medications and hormonal contraceptives that could lower contraceptive efficacy (AII*).• Pediatric and adolescent care providers should prepare adolescents for the transition into adult care settings (AIII).
<p>Rating of Recommendations: A = Strong; B = Moderate; C = Optional</p> <p>Rating of Evidence: I = One or more randomized trials in children† with clinical outcomes and/or validated endpoints; I* = One or more randomized trials in adults with clinical outcomes and/or validated laboratory endpoints with accompanying data in children† from one or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; II = One or more well-designed, nonrandomized trials or observational cohort studies in children† with long-term outcomes; II* = One or more well-designed, nonrandomized trials or observational studies in adults with long-term clinical outcomes with accompanying data in children† from one or more similar nonrandomized trials or cohort studies with clinical outcome data; III = Expert opinion</p> <p>† Studies that include children or children/adolescents, but not studies limited to post-pubertal adolescents.</p>

Background

The majority of individuals in the United States who acquired HIV through perinatal transmission are now adolescents or young adults. Most have had a long clinical course with an extensive antiretroviral (ARV) treatment history.^{1,2} Many older youth and adults may have initially received nonsuppressive monotherapy or dual therapy prior to the availability of combination ARV regimens, including fixed-dose combination (FDC) formulations. Challenges that affect the treatment of adolescents with perinatally acquired HIV include extensive drug resistance, complex regimens, the long-term consequences of HIV and antiretroviral therapy (ART) exposure,³ the developmental transition to adulthood, and psychosocial factors.⁴⁻⁷

In the United States, most adolescents aged ≥ 14 years who recently received HIV diagnoses acquired their infection by horizontal, rather than perinatal, transmission.⁸ They generally follow a clinical course similar to that of adults, and the [Adult and Adolescent Antiretroviral Guidelines](#) should be consulted for treatment recommendations for these patients. Additional information that is specific to the care of post-pubertal adolescents can be found in [Adolescents and Young Adults with HIV](#).

Timing and Selection of Antiretroviral Therapy

All adolescents with HIV (like all people with HIV) should initiate ART as soon as possible **after HIV diagnosis**. Recommendations for ART selection in adolescents with sexual maturity ratings (SMRs) between 1 and 3 can be found in [What to Start](#) and [Appendix A: Pediatric Antiretroviral Drug Information](#). ART recommendations for adolescents and young adults with SMRs between 4 and 5 are available in the [What to Start](#) section of the Adult and Adolescent Antiretroviral Guidelines. Optimizing and simplifying treatment may be especially important when treating adolescents, because this can help improve adherence (see [Modifying Antiretroviral Regimens in Children with Sustained Virologic Suppression on Antiretroviral Therapy](#)). Clinicians who are treating adolescents of childbearing potential should consider additional factors before initiating ART, including potential drug interactions with contraception and the safety of using certain ARV drugs before conception or during pregnancy (see the Contraception, Pregnancy, and Antiretroviral Therapy section below).

Dosing of Antiretroviral Therapy for Adolescents with HIV

Clinical providers need to pay attention to the transition of adolescents from pediatric to adult ART dosing. Many ARV drugs (e.g., abacavir, emtricitabine, lamivudine, tenofovir disoproxil fumarate [TDF], and some protease inhibitors [PIs]) are administered to children at higher body weight–based **doses** or body surface area–based doses than would be predicted by direct extrapolation of adult doses. These doses are based on reported pharmacokinetic data that indicate more rapid drug clearance in children than in adults. Therefore, failure to ensure weight-appropriate dosing in adolescents can result in an increased risk of drug toxicity if higher pediatric dosing is not transitioned to lower adult dosing (often between 25 kg and 40 kg, depending on the particular drug).⁹

Adherence Concerns in Adolescents

Poor adherence to ART is a common problem among adolescents with HIV. Both psychosocial and cognitive developmental factors may contribute to adherence challenges, and these factors should be assessed regularly. The adolescent’s individual needs and preferences also should be considered when making decisions about initiating or changing ART. Comprehensive systems of care are required to serve both the medical and psychosocial needs of adolescents with HIV, because they are frequently inexperienced with managing their health care and may also lack health insurance. Adolescents with perinatally acquired HIV infection are at risk for neurocognitive impairment, which also can interfere with medication adherence.¹⁰ Many also are at risk for mental health comorbidities, including psychiatric, behavioral, and substance use disorders that may interfere with adherence to ART.^{7,11} Compared with adults, youth have lower rates of viral suppression and higher rates of virologic rebound and loss to follow-up.^{12,13} For further discussion of interventions to promote adherence in adolescents, see the [Adolescents and Young Adults with HIV](#) section of the Adult and Adolescent Antiretroviral Guidelines and a 2013 review by Agwu and Fairlie.³

A specific challenge is presented by youth who, despite interventions, remain unable to adhere to therapy. In these cases, alternatives to changing the ARV regimen can include, but are not limited to, simplifying treatment to a once-daily regimen or an FDC tablet; using cellphone alerts and other mHealth approaches to remind patients to take their medication and attend clinic visits; initiating a short-term deferral of treatment until adherence improves or while adherence-related problems (including mental health and substance use disorders) are aggressively addressed; initiating an adherence testing and training period during which a placebo (e.g., vitamin pill) is administered; scheduling appointments more frequently; employing directly observed therapy; and avoiding

regimens with a low genetic resistance threshold. Such decisions should be individualized, and the patient's clinical and laboratory status monitored carefully. The use of long-acting oral and injectable ARV regimens for adolescents is currently being investigated. These regimens may provide an alternative approach for adolescents with adherence challenges.

Mental Health and Substance Use Concerns in Adolescents

Many factors can increase the risk of adverse mental health outcomes among adolescents with perinatally acquired HIV, including long-term medical treatment for a chronic disease, hospitalizations, stigma, the neurocognitive impacts of HIV, parental psychiatric and substance use disorders, and family and caregiver stress and loss. The prevalence of mental health disorders in youth with perinatally acquired HIV is high, with nearly 70% of these adolescents meeting the criteria for a psychiatric disorder at some point in their lives.^{7,14-16} The most common conditions include anxiety and behavioral disorders, mood disorders (including depression), and attention deficit hyperactivity disorder. In at least one cohort, the risk of psychosis and severe chronic mental health disorders was higher in adolescents with perinatally acquired HIV than expected in the general young adult population.¹⁷ Effectively managing psychiatric comorbidities can improve a patient's adherence to medical care, including ART, and can lead to better academic performance and interpersonal relationships (see [Substance Use Disorders and HIV](#) in the Adult and Adolescent Antiretroviral Guidelines).^{11,18-20}

Interventions that address mental health in youth with perinatally acquired HIV include pharmacologic interventions; behavioral modification; and individual, family, and group counseling. The use of telehealth or counseling via videoconferencing may be feasible and acceptable and may improve access to mental health treatment for adolescents with HIV.²¹ However, data are lacking on the effectiveness of these interventions on HIV clinical outcomes.²²⁻²⁴ Current evidence suggests that a combination of tailored psychotherapy—such as cognitive behavioral therapy—and pharmacotherapy can reduce depressive symptoms in adolescents with HIV; however, clinicians who prescribe pharmacotherapy for depression must take potential interactions with ARV drugs into account.^{25,26}

There is evidence that adolescents with perinatally acquired HIV are more likely to have substance use disorders compared to the general population.²⁷ However, available studies on substance use among adolescents with perinatally acquired HIV show age of initiation and rates of substance use similar to age-matched peers without HIV.²⁸ In a comparison of 390 youth with perinatal exposure to HIV versus 211 youth living with perinatally acquired HIV, investigators from the Pediatric HIV/AIDS Cohort Study (PHACS) found that nearly half of both groups had ever used alcohol or marijuana, with a majority having used either substance in the last 3 months, and one out of five marijuana users reporting at least daily use.²⁹ In another study, there was no difference in substance use between adolescents exposed to HIV and adolescents living with HIV. While rates of substance use may not be higher in adolescents with perinatally acquired HIV, the impact on health outcomes—including interference with medication adherence and increased risk taking and decreased safe sex practices—and the potential for comorbid mental health concerns make addressing substance use in adolescents with HIV an important consideration for HIV care providers.^{30,31}

Providers who are caring for adolescents with HIV should incorporate screening for psychiatric and substance use disorders into routine care and refer patients to age-appropriate services as needed. The [American Academy of Pediatrics](#) policy statement provides some guidance and screening tools, particularly for depression. Screening tools for substance use—such as [Screening, Brief Intervention, and Referral to Treatment \(SBIRT\)](#) or [Car, Relax, Alone, Forget, Friends, and Trouble \(CRAFT\)](#)—

may be used.³² Providers also should consider emerging substance use trends when screening adolescents with HIV. Further guidance on screening tools for substance use and mental health is provided by the National Institute on Drug Abuse's [Screening and Assessment Tools Chart](#).

Sexually Transmitted Infections in Adolescents

Clinicians should discuss the risk of sexually transmitted infections (STIs) with their patients. All adolescents with HIV should be screened for STIs and treated appropriately. Clinicians should regularly obtain a detailed sexual history for adolescents to determine which STI screening tests are appropriate. In young men who have sex with men, screening for STIs often requires sampling from several body sites—including the oropharynx, rectum, and urethra—because multiple sites of infection are common. Furthermore, a negative assay at a single site does not exclude infection at another site.³³ For a more detailed discussion of STIs, see the most recent Centers for Disease Control and Prevention guidelines,³⁴ [Human Papillomavirus Disease](#) in the Adult and Adolescent Opportunistic Infection Guidelines, and [Human Papillomavirus](#) in the Pediatric Opportunistic Infection Guidelines. All female adolescents with HIV who are sexually active should receive gynecologic services. All adolescents should receive three doses of the 9-valent human papillomavirus vaccination.

Contraception, Pregnancy, and Antiretroviral Therapy

Adolescents with HIV may initiate sexual activity before or after puberty. Sexually active adolescents are at risk for unintended pregnancy. Approximately half of pregnancies in the United States, including those among women with HIV, are unintended or unplanned.^{35,36} Providers should regularly assess adolescents' desires to become pregnant or avoid pregnancy (also known as their pregnancy intentions). Family planning counseling—including a discussion of the risks of sexual HIV transmission, perinatal HIV transmission, and methods for reducing these risks—should be provided to all youth. Reproductive health options—such as pregnancy planning, preconception care, contraceptive methods, pre-exposure prophylaxis for partners, the concept of Undetectable = Untransmittable (U=U),^{37,38} and safer sex techniques (including instruction on the correct and consistent use of condoms) for prevention of sexual HIV transmission—should be discussed regularly (see [U.S. Medical Eligibility Criteria for Contraceptive Use](#)). For additional information, refer to the following sections of the Perinatal Guidelines: [Prepregnancy Counseling and Care for Persons of Childbearing Age with HIV](#) and [Reproductive Options When One or Both Partners Have HIV](#). The American Academy of Pediatrics Committee on Adolescence offers guidance about the integration of sexual and reproductive health care in pediatric clinical settings.³⁹

The possibility of planned and unplanned pregnancy should be considered when selecting an ARV regimen for a female adolescent. The most vulnerable period in fetal organogenesis is the first trimester, often before pregnancy is recognized. When treating adolescents of childbearing potential, clinicians should carefully review the potential toxicities of ARV drugs and consider making any necessary changes to a regimen as promptly as possible (e.g., before conception, when possible). For information about the selection and management of ARV drugs before and during pregnancy for people with HIV who are of childbearing age, see [Table 5](#) in the [Recommendations for Use of Antiretroviral Drugs During Pregnancy](#) section of the Perinatal Guidelines. When discussing ART options with female adolescents and their caregivers, it is important to consider the benefits and risks of all ARV drugs and to provide the information and counseling needed to support informed decision-making (see [Appendix C: Antiretroviral Counseling Guide for Health Care Providers](#)).

Interactions Between Contraceptives and Antiretroviral Drugs

People living with HIV can use all available contraceptive methods, including hormonal contraceptives, implantable devices, intrauterine devices, the transdermal patch, and a vaginal ring.⁴⁰

Several PIs and non-nucleoside reverse transcriptase inhibitors alter the metabolism of oral contraceptives, which theoretically may reduce the efficacy of oral contraceptive agents or increase the risk of estrogen-related or progestin-related adverse effects⁴¹⁻⁴³ (see [Drug–Drug Interactions](#) in the [Adult and Adolescent Antiretroviral Guidelines](#) and the [HIV Drug Interaction Checker](#)). Integrase strand transfer inhibitors appear to have no interaction with estrogen-based contraceptives.^{44,45} For more information about potential interactions between ARV drugs and hormonal contraceptives, see [Table 3](#) in the Perinatal Guidelines.

Concerns about loss of bone mineral density with long-term use of depot medroxyprogesterone acetate (DMPA), with or without coadministration of ART (specifically TDF), should not preclude the use of DMPA as an effective contraceptive, unless clinical evidence indicates bone fragility.

Pregnant Adolescents with HIV

Adolescents who want to become pregnant should receive preconception counseling and care, including a discussion of pregnancy planning and special considerations when using ARV drugs during pregnancy (see [Prepregnancy Counseling and Care for Persons of Childbearing Age with HIV](#) in the Perinatal Guidelines). Pregnancy should not preclude the use of optimal therapeutic ARV regimens. Clinicians need to consider maternal and fetal safety, as well as the need to prevent perinatal transmission of HIV, when selecting regimens for pregnant people or adolescents who are planning to become pregnant. See the [Prepregnancy Counseling and Care for Persons of Childbearing Age with HIV](#) for more details about choosing an ARV regimen for pregnant people with HIV, including adolescents. Pregnancies occur as adolescents with perinatally acquired HIV enter adolescence and young adulthood.^{46,47} Some studies suggest higher rates of adverse pregnancy outcomes—such as small-for-gestational-age infants—among pregnant people with perinatally acquired HIV than among those who acquired HIV by horizontal transmission. Unplanned pregnancy is not uncommon in youth living with perinatally acquired HIV.⁴⁷⁻⁴⁹ Women with perinatally acquired HIV also may be more likely to have complex ARV histories, virologic failure, and drug resistance at the time of pregnancy.⁵⁰⁻⁵² However, the rate of perinatal transmission among pregnant people with perinatally acquired HIV who are receiving ART appears to be similar to the rate among people on ART who acquired HIV by horizontal transmission.⁵³⁻⁵⁷

Special Considerations for Adolescents with HIV Who Are Sexual Minorities

Adolescence represents a period of emerging recognition of sexual identity. Adolescents **with HIV** who are lesbian, gay, bisexual, transgender, or nonbinary require both culturally competent providers and tailored medical care. Health care providers should ask patients nonjudgmental questions about their sexual and gender identity to determine whether they require specific medical and support services. It is important to consider the possibility of drug–drug interactions in adolescents who are receiving both ART and gender-affirming hormone therapy. Additional resources for the care of these adolescents can be found in the [Adolescents and Young Adults with HIV](#) section and the [Transgender People with HIV](#) section of the [Adult and Adolescent Antiretroviral Guidelines](#).

Transitioning Adolescents into Adult HIV Care Settings

Transition, as defined by Reiss et. al., may be described as “a multifaceted, active process that attends to the medical, psychosocial, cognitive and educational, or vocational needs of adolescents as they move from the child- to the adult-focused health care system.”⁵⁸ Facilitating a successful transition for adolescents with HIV from their pediatric/adolescent care clinic to adult care is important, but challenging.⁵⁹⁻⁶² Many adolescents disengage from care during the transition to adult care, putting them at risk for HIV progression and transmission to partners.⁶³⁻⁶⁵ Pediatric and adolescent care providers and their multidisciplinary teams should have a formal written plan in place to transition adolescents to adult care. Although transition generally occurs when individuals are in their late teens or early 20s, discussion of and planning for the transition process should be initiated early in the teen years, with involvement from both the adolescent and their parents and/or caregivers. Care models for children and adolescents with perinatally acquired HIV tend to be family centered, consisting of a multidisciplinary team that often includes physicians, nurses, social workers, and mental health professionals. These providers generally have long-standing relationships with patients and their families, and care is rendered in discreet, intimate settings. Although expert care also is provided under the adult HIV care medical model, adolescents and their caregivers may be unfamiliar with the busier, more individual-centered clinics that are typical of adult medical care providers. These providers often expect patients to assume a greater level of responsibility for their care, and adolescents may be uncomfortable with providers with whom they do not have a long-standing relationship.

One multisite study in the United States found that adolescents who transitioned to adult care at an older age reported greater satisfaction with their care than those who transitioned at a younger age. Additionally, adolescents who reported being able to perform certain tasks that were related to their care (e.g., making appointments, requesting prescriptions, arranging transportation to appointments) were more likely to be engaged in adult care.⁶⁶ It may be beneficial to provide adolescents, caregivers, and their new adult medical care providers with support and guidance regarding the expectations for each person involved in the patient–provider relationship. In this situation, it may be helpful for a pediatric care provider and an adult care provider to share joint care of a patient for a period.

Adolescent care providers should have a candid discussion with the transitioning adolescent and their caregivers to understand what qualities the adolescent considers most important when choosing an adult care setting (e.g., confidentiality, small clinic size, low patient-to-provider ratio, availability of after-school or evening appointments). Social determinants—such as the patient’s developmental status, behavioral/mental health comorbidities, housing, family support, employment status, recent discharge from foster care, peer pressure, illicit drug use, and incarceration—should be considered during transition.

No definitive model of transition to adult HIV care currently exists, and only a limited number of studies have reported on outcomes following transition, although research in this area is ongoing. However, emerging qualitative research has revealed the importance of the patient–provider relationship, including trust, the need for developmentally appropriate preparation for transition, and opportunities for growth and independence.^{62,67} Recent studies have shown potential for successful transition and ongoing retention using models that include a multidisciplinary approach, which utilizes providers co-trained in both internal medicine and pediatrics, peer navigators, social workers, mental health support, and a youth-focused care model for adolescents who were already attending adult HIV clinics.^{68,69}

Several studies have shown that youth with HIV who transitioned into adult care settings had higher rates of attrition from care than those who remained in pediatric/adolescent care; U.S. studies show that less than half of youth who transitioned care to an adult clinic remained in care after 9 to 12 months.^{63,64,70} In addition to poor retention in care, several studies have identified poor viral suppression rates in transitioned youth and young adults with HIV.² Pre-transition virologic failure and longer linkage times have been associated with worse outcomes post-transition.^{62,65} Furthermore, some reports from the United Kingdom suggest that the mortality rate of adolescents with HIV increases after transition,^{19,65,71} underscoring the need to critically examine transition and determine the best mechanisms to optimize the long-term outcomes for youth with perinatally acquired HIV.⁶³

Some general guidelines, mostly based on anecdotal evidence and consensus expert opinion, are available about transition plans and who might benefit most from them.^{60,72-79} To maximize the likelihood of success, providers should prepare adolescents for transition long before it occurs. Attention to the following key areas could improve retention in care and minimize the risk of ART interruptions:

- Educating HIV care teams and staff about transitioning;
- Beginning discussions about transition early, before the actual transition process;
- Developing a written, individualized transition plan to address comprehensive care needs, including medical, psychosocial, and financial aspects of transitioning;
- Optimizing communication between providers at pediatric/adolescent clinics and providers at adult clinics;
- Identifying adult care providers who are experts in providing care to adolescents and young adults;
- Fostering a trusting patient–provider relationship with new adult care providers;
- Addressing barriers caused by a lack of information, stigma, or disclosure concerns;
- Discussing the differences between the practice styles of adult clinics and pediatric/adolescent clinics;
- Helping youth develop the skills needed to manage their care, including counseling them on appointment management, the appropriate use of a primary care provider, the importance of prompt symptom recognition and reporting, and the importance of managing medications, insurance, and state and federal benefits;
- Identifying an optimal clinic model for a given setting (e.g., simultaneous transition of mental health and/or case management services versus a gradual phase-in);
- Clearly defining the desired outcomes for the transition, such as retention in care, ongoing access to other services (e.g., case management, mental health), clinical outcomes (e.g., viral suppression), and patient satisfaction;
- Implementing ongoing evaluations to measure the success of a transition model;
- Engaging in regular multidisciplinary case conferences between adult and adolescent care providers;
- Implementing interventions that may be associated with improved outcomes, such as support groups and mental health consultation; *and*
- Identifying a care navigator who can provide support during the transition.

References

1. Van Dyke RB, Patel K, Siberry GK, et al. Antiretroviral treatment of US children with perinatally acquired HIV infection: temporal changes in therapy between 1991 and 2009 and predictors of immunologic and virologic outcomes. *J Acquir Immune Defic Syndr*. 2011;57(2):165-173. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21407086>.
2. Xia Q, Shah D, Gill B, Torian LV, Braunstein SL. Continuum of care among people living with perinatally acquired HIV infection in New York City, 2014. *Public Health Rep*. 2016;131(4):566-573. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27453601>.
3. Agwu AL, Fairlie L. Antiretroviral treatment, management challenges and outcomes in perinatally HIV-infected adolescents. *J Int AIDS Soc*. 2013;16:18579. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23782477>.
4. Judd A, Lodwick R, Noguera-Julian A, et al. Higher rates of triple-class virological failure in perinatally HIV-infected teenagers compared with heterosexually infected young adults in Europe. *HIV Med*. 2017;18(3):171-180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27625109>.
5. Hermetet-Lindsay KD, Correia KF, Williams PL, et al. Contributions of disease severity, psychosocial factors, and cognition to behavioral functioning in U.S. youth perinatally exposed to HIV. *AIDS Behav*. 2017;21(9):2703-2715. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27475941>.
6. Tarantino N, Brown LK, Whiteley L, et al. Correlates of missed clinic visits among youth living with HIV. *AIDS Care*. 2018;30(8):982-989. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29455553>.
7. Bucek A, Leu CS, Benson S, et al. Psychiatric disorders, antiretroviral medication adherence and viremia in a cohort of perinatally HIV-infected adolescents and young adults. *Pediatr Infect Dis J*. 2018;37(7):673-677. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29227462>.
8. Centers for Disease Control and Prevention. HIV surveillance supplemental report, 2018 (updated). 2020. Available at: <https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>.
9. Rakhmanina N, Phelps BR. Pharmacotherapy of pediatric HIV infection. *Pediatr Clin North Am*. 2012;59(5):1093-1115. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23036246>.
10. Harris LL, Chernoff MC, Nichols SL, et al. Prospective memory in youth with perinatally-acquired HIV infection. *Child Neuropsychol*. 2018;24(7):938-958. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28782457>.
11. Mellins CA, Tassiopoulos K, Malee K, et al. Behavioral health risks in perinatally HIV-exposed youth: co-occurrence of sexual and drug use behavior, mental health problems,

- and nonadherence to antiretroviral treatment. *AIDS Patient Care STDS*. 2011;25(7):413-422. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21992620>.
12. Gray KM, Ocfemia MCB, Wang X, Li J, Nesheim SR. Characteristics and care outcomes among persons living with perinatally acquired HIV infection in the United States, 2015. *J Acquir Immune Defic Syndr*. 2019;82(1):17-23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31169773>.
 13. Kacanek D, Huo Y, Malee K, et al. Nonadherence and unsuppressed viral load across adolescence among U.S. youth with perinatally acquired HIV. *AIDS*. 2019;33(12):1923-1934. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31274538>.
 14. Mellins CA, Malee KM. Understanding the mental health of youth living with perinatal HIV infection: lessons learned and current challenges. *J Int AIDS Soc*. 2013;16:18593. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23782478>.
 15. Mellins CA, Brackis-Cott E, Leu CS, et al. Rates and types of psychiatric disorders in perinatally human immunodeficiency virus–infected youth and seroreverters. *J Child Psychol Psychiatry*. 2009;50(9):1131-1138. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19298479>.
 16. Gadow KD, Chernoff M, Williams PL, et al. Co-occurring psychiatric symptoms in children perinatally infected with HIV and peer comparison sample. *J Dev Behav Pediatr*. 2010;31(2):116-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20110828>.
 17. Mallik I, Pasvol T, Frize G, et al. Psychotic disorders in young adults with perinatally acquired HIV: a U.K. case series. *Psychol Med*. 2020:1-7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33183361>.
 18. Kapetanovic S, Wiegand RE, Dominguez K, et al. Associations of medically documented psychiatric diagnoses and risky health behaviors in highly active antiretroviral therapy–experienced perinatally HIV-infected youth. *AIDS Patient Care STDS*. 2011;25(8):493-501. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21745118>.
 19. Fish R, Judd A, Jungmann E, O’Leary C, Foster C, HIV Young Persons Network. Mortality in perinatally HIV-infected young people in England following transition to adult care: an HIV Young Persons Network (HYPNet) audit. *HIV Med*. 2013;15(4):239-244. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24112550>.
 20. Kim SH, Gerver SM, Fidler S, Ward H. Adherence to antiretroviral therapy in adolescents living with HIV: systematic review and meta-analysis. *AIDS*. 2014;28(13):1945-1956. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24845154>.
 21. Saberi P, McCuistian C, Agnew E, et al. Video-counseling intervention to address HIV care engagement, mental health, and substance use challenges: a pilot randomized clinical trial for youth and young adults living with HIV. *Telemed Rep*. 2021;2(1):14-25. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33575683>.

22. Chernoff M, Nachman S, Williams P, et al. Mental health treatment patterns in perinatally HIV-infected youth and controls. *Pediatrics*. 2009;124(2):627-636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19596734>.
23. Sirois PA, Montepiedra G, Kapetanovic S, et al. Impact of medications prescribed for treatment of attention-deficit hyperactivity disorder on physical growth in children and adolescents with HIV. *J Dev Behav Pediatr*. 2009;30(5):403-412. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19827220>.
24. Funck-Brentano I, Dalban C, Veber F, et al. Evaluation of a peer support group therapy for HIV-infected adolescents. *AIDS*. 2005;19(14):1501-1508. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16135904>.
25. Brown LK, Kennard BD, Emslie GJ, et al. Effective treatment of depressive disorders in medical clinics for adolescents and young adults living with HIV: a controlled trial. *J Acquir Immune Defic Syndr*. 2016;71(1):38-46. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26761270>.
26. Benton TD, Kee Ng WY, Leung D, Canetti A, Karnik N. Depression among youth living with HIV/AIDS. *Child Adolesc Psychiatr Clin N Am*. 2019;28(3):447-459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31076119>.
27. Abrams EJ, Mellins CA, Bucek A, et al. Behavioral health and adult milestones in young adults with perinatal HIV infection or exposure. *Pediatrics*. 2018;142(3). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30097528>.
28. Alperen J, Brummel S, Tassiopoulos K, et al. Prevalence of and risk factors for substance use among perinatally human immunodeficiency virus–infected and perinatally exposed but uninfected youth. *J Adolesc Health*. 2014;54(3):341-349. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24239286>.
29. Nichols SL, Brummel S, Malee KM, et al. The role of behavioral and neurocognitive functioning in substance use among youth with perinatally acquired HIV infection and perinatal HIV exposure without infection. *AIDS Behav*. 2021;25(9):2827-2840. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33616833>.
30. Williams PL, Leister E, Chernoff M, et al. Substance use and its association with psychiatric symptoms in perinatally HIV-infected and HIV-affected adolescents. *AIDS Behav*. 2010;14(5):1072-1082. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20725774>.
31. Elkington KS, Cruz JE, Warne P, Santamaria EK, Dolezal C, Mellins CA. Marijuana use and psychiatric disorders in perinatally HIV-exposed youth: does HIV matter? *J Pediatr Psychol*. 2016;41(3):277-286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26698841>.
32. Knight JR, Shrier LA, Bravender TD, Farrell M, Vander Bilt J, Shaffer HJ. A new brief screen for adolescent substance abuse. *Arch Pediatr Adolesc Med*. 1999;153(6):591-596. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10357299>.

33. Sultan B, White JA, Fish R, et al. The ‘3 in 1’ study: pooling self-taken pharyngeal, urethral and rectal samples into a single sample for analysis, for diagnosis of *Neisseria gonorrhoeae* and *Chlamydia trachomatis* in men who have sex with men (MSM). *J Clin Microbiol*. 2015;54(3):650-656. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26719439>.
34. Workowski KA, Bolan GA, Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines, 2015. *MMWR Recomm Rep*. 2015;64(RR-03):1-137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26042815>.
35. Finer LB, Zolna MR. Declines in unintended pregnancy in the United States, 2008–2011. *N Engl J Med*. 2016;374(9):843-852. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26962904>.
36. Sutton MY, Patel R, Frazier EL. Unplanned pregnancies among HIV-infected women in care—United States. *J Acquir Immune Defic Syndr*. 2014;65(3):350-358. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24189153>.
37. Cohen MS, Chen YQ, McCauley M, et al. Antiretroviral therapy for the prevention of HIV-1 transmission. *N Engl J Med*. 2016;375(9):830-839. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27424812>.
38. Centers for Disease Control and Prevention. Evidence of HIV treatment and viral suppression in preventing the sexual transmission of HIV. 2018. Available at: <https://www.cdc.gov/hiv/pdf/risk/art/cdc-hiv-art-viral-suppression.pdf>.
39. Marcell AV, Burstein GR, Committee on Adolescence. Sexual and reproductive health care services in the pediatric setting. *Pediatrics*. 2017;140(5). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29061870>.
40. Panel on Treatment of HIV During Pregnancy and Prevention of Perinatal Transmission. Recommendations for the use of antiretroviral drugs during pregnancy and interventions to reduce perinatal HIV transmission in the United States. 2021. Available at: <https://clinicalinfo.hiv.gov/en/guidelines/perinatal/guidelines-panel-members?view=full>.
41. Sevinsky H, Eley T, Persson A, et al. The effect of efavirenz on the pharmacokinetics of an oral contraceptive containing ethinyl estradiol and norgestimate in healthy HIV-negative women. *Antivir Ther*. 2011;16(2):149-156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21447863>.
42. Zhang J, Chung E, Yones C, et al. The effect of atazanavir/ritonavir on the pharmacokinetics of an oral contraceptive containing ethinyl estradiol and norgestimate in healthy women. *Antivir Ther*. 2011;16(2):157-164. Available at: <https://ncbi.nlm.nih.gov/pubmed/21447864>.
43. El-Ibiary SY, Cocohoba JM. Effects of HIV antiretrovirals on the pharmacokinetics of hormonal contraceptives. *Eur J Contracept Reprod Health Care*. 2008;13(2):123-132. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18465473>.

44. Song IH, Borland J, Chen S, Wajima T, Peppercorn AF, Piscitelli SC. Dolutegravir has no effect on the pharmacokinetics of oral contraceptives with norgestimate and ethinyl estradiol. *Ann Pharmacother*. 2015;49(7):784-789. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25862012>.
45. Patel R, Stalter R, Onono M, et al. Dolutegravir-containing ART does not reduce etonogestrel implant concentrations. Abstract 129. Presented at: Conference on Retroviruses and Opportunistic Infections 2020. Boston, MA. Available at: <https://www.croiconference.org/abstract/dolutegravir-containing-art-does-not-reduce-etonogestrel-implant-concentrations>.
46. Kenny J, Williams B, Prime K, Tookey P, Foster C. Pregnancy outcomes in adolescents in the U.K. and Ireland growing up with HIV. *HIV Med*. 2012;13(5):304-308. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22136754>.
47. Byrne L, Thorne C, Foster C, Tookey P. Pregnancy outcomes in women growing up with perinatally acquired HIV in the United Kingdom and Ireland. *J Int AIDS Soc*. 2014;17(4 Suppl 3):19693. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25397443>.
48. Jao J, Agwu A, Mhango G, et al. Growth patterns in the first year of life differ in infants born to perinatally vs. nonperinatally HIV-infected women. *AIDS*. 2015;29(1):111-116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25562495>.
49. Jao J, Sigel KM, Chen KT, et al. Small for gestational age birth outcomes in pregnant women with perinatally acquired HIV. *AIDS*. 2012;26(7):855-859. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22313958>.
50. Lazenby GB, Mmeje O, Fisher BM, et al. Antiretroviral resistance and pregnancy characteristics of women with perinatal and nonperinatal HIV infection. *Infect Dis Obstet Gynecol*. 2016;2016:4897501. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27413359>.
51. Cruz ML, Santos E, Benamor Teixeira Mde L, et al. Viral suppression and resistance in a cohort of perinatally-HIV infected (PHIV+) pregnant women. *Int J Environ Res Public Health*. 2016;13(6). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27338425>.
52. Murphy E, Keller J, Argani C, et al. Pregnancy in an urban cohort of adolescents living with human immunodeficiency virus: characteristics and outcomes in comparison to adults. *AIDS Patient Care STDS*. 2021;35(4):103-109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33835849>.
53. Meloni A, Tuveri M, Floridia M, et al. Pregnancy care in two adolescents perinatally infected with HIV. *AIDS Care*. 2009;21(6):796-798. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19806493>.
54. Williams SF, Keane-Tarchichi MH, Bettica L, Dieudonne A, Bardeguez AD. Pregnancy outcomes in young women with perinatally acquired human immunodeficiency virus-1. *Am J Obstet Gynecol*. 2009;200(2):149 e141-145. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18973871>.

55. Cruz ML, Cardoso CA, Joao EC, et al. Pregnancy in HIV vertically infected adolescents and young women: a new generation of HIV-exposed infants. *AIDS*. 2010;24(17):2727-2731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20827164>.
56. Elgalib A, Hegazi A, Samarawickrama A, et al. Pregnancy in HIV-infected teenagers in London. *HIV Med*. 2011;12(2):118-123. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20807252>.
57. Calitri C, Gabiano C, Galli L, et al. The second generation of HIV-1 vertically exposed infants: a case series from the Italian register for paediatric HIV infection. *BMC Infect Dis*. 2014;14:277. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24885649>.
58. Reiss JG, Gibson RW, Walker LR. Health care transition: youth, family, and provider perspectives. *Pediatrics*. 2005;115(1):112-120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15629990>.
59. Tepper V, Zaner S, Ryscavage P. HIV healthcare transition outcomes among youth in North America and Europe: a review. *J Int AIDS Soc*. 2017;20(Suppl 3):60-70. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28530041>.
60. Judd A, Davies MA. Adolescent transition among young people with perinatal HIV in high-income and low-income settings. *Curr Opin HIV AIDS*. 2018;13(3):236-248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29528851>.
61. Foster C, Fidler S. Optimizing HIV transition services for young adults. *Curr Opin Infect Dis*. 2018;31(1):33-38. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29210712>.
62. Ritchwood TD, Malo V, Jones C, et al. Healthcare retention and clinical outcomes among adolescents living with HIV after transition from pediatric to adult care: a systematic review. *BMC Public Health*. 2020;20(1):1195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32746881>.
63. Ryscavage P, Macharia T, Patel D, Palmeiro R, Tepper V. Linkage to and retention in care following healthcare transition from pediatric to adult HIV care. *AIDS Care*. 2016;28(5):561-565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26766017>.
64. Tanner AE, Philbin MM, Chambers BD, et al. Healthcare transition for youth living with HIV: outcomes from a prospective multi-site study. *J Adolesc Health*. 2018;63(2):157-165. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29887488>.
65. Hussen SA, Chakraborty R, Knezevic A, et al. Transitioning young adults from paediatric to adult care and the HIV care continuum in Atlanta, Georgia, USA: a retrospective cohort study. *J Int AIDS Soc*. 2017;20(1):1-9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28872281>.
66. Tassiopoulos K, Huo Y, Patel K, et al. Healthcare transition outcomes among young adults with perinatally acquired human immunodeficiency virus infection in the United States. *Clin Infect Dis*. 2020;71(1):133-141. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31584617>.

67. Barr EA, Raybin JL, Dunlevy H, Abuogi L, Jones J. Transition from pediatric and adolescent HIV care to adult HIV care and the patient-provider relationship: a qualitative metasynthesis. *J Assoc Nurses AIDS Care*. 2021. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33654006>.
68. Griffith D, Jin L, Childs J, Posada R, Jao J, Agwu A. Outcomes of a comprehensive retention strategy for youth with HIV after transfer to adult care in the United States. *Pediatr Infect Dis J*. 2019;38(7):722-726. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30985513>.
69. Griffith D, Snyder J, Dell S, Nolan K, Keruly J, Agwu A. Impact of a youth-focused care model on retention and virologic suppression among young adults with HIV cared for in an adult HIV clinic. *J Acquir Immune Defic Syndr*. 2019;80(2):e41-e47. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30422910>.
70. Farmer C, Yehia BR, Fleishman JA, et al. Factors associated with retention among non-perinatally HIV-infected youth in the HIV research network. *J Pediatric Infect Dis Soc*. 2016;5(1):39-46. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26908490>.
71. Foster C, Ayers S, McDonald S, et al. Clinical outcomes post transition to adult services in young adults with perinatally acquired HIV infection: mortality, retention in care and viral suppression. *AIDS*. 2020;34(2):261-266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31651427>.
72. Rosen DS, Blum RW, Britto M, Sawyer SM, Siegel DM, Society for Adolescent Medicine. Transition to adult health care for adolescents and young adults with chronic conditions: position paper of the Society for Adolescent Medicine. *J Adolesc Health*. 2003;33(4):309-311. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14519573>.
73. Gilliam PP, Ellen JM, Leonard L, Kinsman S, Jevitt CM, Straub DM. Transition of adolescents with HIV to adult care: characteristics and current practices of the Adolescent Trials Network for HIV/AIDS interventions. *J Assoc Nurses AIDS Care*. 2011;22(4):283-294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20541443>.
74. New York State Department of Health AIDS Institute. Transitioning HIV-infected adolescents into adult care. 2011. Available at: https://www.medscape.com/viewarticle/748356_2.
75. Andiman WA. Transition from pediatric to adult healthcare services for young adults with chronic illnesses: the special case of human immunodeficiency virus infection. *J Pediatr*. 2011;159(5):714-719. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21868035>.
76. Dowshen N, D'Angelo L. Health care transition for youth living with HIV/AIDS. *Pediatrics*. 2011;128(4):762-771. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21930548>.
77. Committee On Pediatric AIDS. Transitioning HIV-infected youth into adult health care. *Pediatrics*. 2013;132(1):192-197. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23796739>.

78. Sharer M, Fullem A. Transitioning of care and other services for adolescents living with HIV in Sub-Saharan Africa. Arlington, VA: USAID's AIDS Support and Technical Assistance Resources, AIDSTAR-One, Task Order 1.; 2012. Available at: <https://www.socialserviceworkforce.org/resources/transitioning-care-and-other-services-adolescents-living-hiv-sub-saharan-africa>
79. Hussen SA, Chahroudi A, Boylan A, Camacho-Gonzalez AF, Hackett S, Chakraborty R. Transition of youth living with HIV from pediatric to adult-oriented healthcare: a review of the literature. *Future Virol.* 2015;9(10):921-929. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25983853>.

