Appendix D: Supplemental Information

Updated: Apr.11, 2022 Reviewed: Apr.11, 2022

Table A. Likelihood of Developing AIDS or Death Within 12 Months, by Age and CD4 T-Cell Percentage or Log₁₀ HIV-1 RNA Copy Number in HIV-Infected Children Receiving No Therapy or Zidovudine Monotherapy

CD4 Percentage				Log ₁₀ HIV RNA Copy Number			
Age	10%	20%	25%	30%	6.0	5.0	4.0
Percent Mortality (95% Confidence Interval)							
6 Months	28.7	12.4	8.5	6.4	9.7	4.1	2.7
1 Year	19.5	6.8	4.5	3.3	8.8	3.1	1.7
2 Years	11.7	3.1	2.0	1.5	8.2	2.5	1.1
5 Years	4.9	0.9	0.6	0.5	7.8	2.1	0.7
10 Years	2.1	0.3	0.2	0.2	7.7	2.0	0.6
Percent Developing AIDS (95% Confidence Interval)							
6 Months	51.4	31.2	24.9	20.5	23.7	13.6	10.9
1 Year	40.5	20.9	15.9	12.8	20.9	10.5	7.8
2 Years	28.6	12.0	8.8	7.2	18.8	8.1	5.3
5 Years	14.7	4.7	3.7	3.1	17.0	6.0	3.2
10 Years	7.4	2.2	1.9	1.8	16.2	5.1	2.2

Note: Table modified from: HIV Paediatric Prognostic Markers Collaborative Study Group. Lancet. 2003;362:1605-1611.

Table B. Death and AIDS/Death Rate per 100 Person-Years by Current Absolute CD4 Cell Count andAge in HIV-Infected Children Receiving No Therapy or Zidovudine Monotherapy (HIV Paediatric Prognostic Markers Collaborative Study) and Adult Seroconverters (CASCADE Study)

Age (Years)	Absolute CD4 Cell Count (cells/mm ³)						
	<50	50–99	100–199	200–349	350-499	500+	
Rate of Death Per 100 Patient-Years							
0–4	59.3	39.6	25.4	11.1	10.0	3.5	
5–14	28.9	11.8	4.3	0.89	0.00	0.00	
15–24	34.7	6.1	1.1	0.71	0.58	0.65	
25–34	47.7	10.8	3.7	1.1	0.38	0.22	
35–44	58.8	15.6	4.5	0.92	0.74	0.85	
45–54	66.0	18.8	7.7	1.8	1.3	0.86	
55+	91.3	21.4	17.6	3.8	2.5	0.91	

Rate of AIDS or Death per 100 Patient-Years						
0–4	82.4	83.2	57.3	21.4	20.7	14.5
5–14	64.3	19.6	16.0	6.1	4.4	3.5
15–24	61.7	30.2	5.9	2.6	1.8	1.2
25–34	93.2	57.6	19.3	6.1	2.3	1.1
35-44	88.1	58.7	25.5	6.6	4.0	1.9
45–54	129.1	56.2	24.7	7.7	3.1	2.7
55+	157.9	42.5	30.0	10.0	5.1	1.8

Note: Table modified from: HIV Paediatric Prognostic Markers Collaborative Study and the CASCADE Collaboration. *J Infect Dis.* 2008;197:398-404.

Table C. Association of Baseline Human Immunodeficiency Virus (HIV) RNA CopyNumber and CD4T-Cell Percentage with Long-Term Risk of Death in HIV-InfectedChildren^a

Baseline HIV RNA ^c (Copies/mL)	No. Patients ^d	Deat	Deaths ^b			
Baseline CD4 Percentage		Number	Percentage			
≤100,000						
≥15%	103	15	(15%)			
<15%	24	15	(63%)			
>100,000						
≥15%	89	32	(36%)			
<15%	36	29	(81%)			

^a Data from the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Intravenous Immunoglobulin Clinical Trial.

^b Mean follow-up: 5.1 years.

^c Tested by NASBA[®] assay (manufactured by Organon Teknika, Durham, North Carolina) on frozen stored serum.

^d Mean age: 3.4 years.

Source: Mofenson LM, Korelitz J, Meyer WA, et al. The relationship between serum human immunodeficiency virus type 1 (HIV-1) RNA level, CD4 lymphocyte percent, and long-term mortality risk in HIV-1-infected children. J Infect Dis. 1997;175(5):1029–1038.

Figure A. Estimated Probability of AIDS Within 12 Months by Age and CD4 Percentage in HIV-Infected Children Receiving No Therapy or Zidovudine Monotherapy





Figure modified from Lancet 2003;362:1605-1611






Figure modified from Lancet 2003;362:1605-1611

Figure C. Death Rate per 100 Person-Years in HIV-Infected Children Aged 5 Years or Older in the HIV Paediatric Prognostic Marker Collaborative Study and HIV-Infected Seroconverting Adults from the CASCADE Study*




Figure modifed from: HIV Paediatric Prognostic Markers Collaborative Study and the CASCADE Collaboration. *J Infect Dis.* 2008;197:398-404.

Figure D. Estimated Probability of AIDS Within 12 Months of Age and HIV RNA Copy Number in HIV-Infected Children Receiving No Therapy or Zidovudine Monotherapy




Figure modified from Lancet 2003;362:1605-1611



